


**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

|   |  |   |
|---|--|---|
| <b>(51) International Patent Classification <sup>6</sup> :</b><br><b>C07K 14/47, 16/18, C12N 15/12</b>  | <b>A2</b>  | <b>(11) International Publication Number:</b> <b>WO 95/32221</b><br><b>(43) International Publication Date:</b> 30 November 1995 (30.11.95) |
| <b>(21) International Application Number:</b> PCT/US95/06613<br><b>(22) International Filing Date:</b> 24 May 1995 (24.05.95)<br><br><b>(30) Priority Data:</b><br>08/249,182 25 May 1994 (25.05.94) US<br>08/346,455 28 November 1994 (28.11.94) US<br><br><b>(71) Applicant:</b> THE GOVERNMENT OF THE UNITED STATES OF AMERICA, represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; Office of Technology Transfer, National Institutes of Health, Box 13, Suite 325, 6011 Executive Boulevard, Rockville, MD 20852 (US).<br><br><b>(72) Inventors:</b> STRACKE, Mary; 14414 Barkwood Drive, Rockville, MD 20853 (US). LIOTTA, Lance; 9027 Mistwood Drive, Potomac, MD 20854 (US). SCHIFFMANN, Elliott; 3027 Pickwick Lane, Chevy Chase, MD 20815 (US). KRUTZCH, Jerry; 9704 De Paul Drive, Bethesda, MD 20817 (US). MURATA, Jun; Dept. of Pathogenic Biochemistry, Research Institute of Wakan-Yaku, Toyoma Medical & Pharmaceutical University, 2630 Sugitani, Toyama 930-01 (JP). | <b>(74) Agents:</b> FEILER, William, S. et al.; Morgan & Finnegan, L.L.P., 345 Park Avenue, New York, NY 10154 (US).<br><br><b>(81) Designated States:</b> AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).<br><br><b>Published</b><br><i>Without international search report and to be republished upon receipt of that report.</i> |   |

**(54) Title:** AUTOTAXIN: MOTILITY STIMULATING PROTEIN USEFUL IN CANCER DIAGNOSIS AND THERAPY**(57) Abstract**

The present invention relates, in general, to autotaxin. In particular, the present invention relates to a DNA segment encoding autotaxin; recombinant DNA molecules containing the DNA segment; cells containing the recombinant DNA molecule; a method of producing autotaxin; antibodies to autotaxin; and identification of functional domains in autotaxin.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

|    |                          |    |                                       |    |                          |
|----|--------------------------|----|---------------------------------------|----|--------------------------|
| AT | Austria                  | GB | United Kingdom                        | MR | Mauritania               |
| AU | Australia                | GE | Georgia                               | MW | Malawi                   |
| BB | Barbados                 | GN | Guinea                                | NE | Niger                    |
| BE | Belgium                  | GR | Greece                                | NL | Netherlands              |
| BF | Burkina Faso             | HU | Hungary                               | NO | Norway                   |
| BG | Bulgaria                 | IE | Ireland                               | NZ | New Zealand              |
| BJ | Benin                    | IT | Italy                                 | PL | Poland                   |
| BR | Brazil                   | JP | Japan                                 | PT | Portugal                 |
| BY | Belarus                  | KE | Kenya                                 | RO | Romania                  |
| CA | Canada                   | KG | Kyrgyzstan                            | RU | Russian Federation       |
| CF | Central African Republic | KP | Democratic People's Republic of Korea | SD | Sudan                    |
| CG | Congo                    | KR | Republic of Korea                     | SE | Sweden                   |
| CH | Switzerland              | KZ | Kazakhstan                            | SI | Slovenia                 |
| CI | Côte d'Ivoire            | LI | Liechtenstein                         | SK | Slovakia                 |
| CM | Cameroon                 | LK | Sri Lanka                             | SN | Senegal                  |
| CN | China                    | LU | Luxembourg                            | TD | Chad                     |
| CS | Czechoslovakia           | LV | Latvia                                | TG | Togo                     |
| CZ | Czech Republic           | MC | Monaco                                | TJ | Tajikistan               |
| DE | Germany                  | MD | Republic of Moldova                   | TT | Trinidad and Tobago      |
| DK | Denmark                  | MG | Madagascar                            | UA | Ukraine                  |
| ES | Spain                    | ML | Mali                                  | US | United States of America |
| FI | Finland                  | MN | Mongolia                              | UZ | Uzbekistan               |
| FR | France                   |    |                                       | VN | Viet Nam                 |
| GA | Gabon                    |    |                                       |    |                          |

- 1 -

AUTOTAXIN: MOTILITY STIMULATING PROTEIN USEFUL  
IN CANCER DIAGNOSIS AND THERAPY

This application is a continuation-in-part of  
application serial no. 08/249,182 filed May 25, 1994,  
which is a continuation-in-part of application serial no.  
07/822,043 filed on Jan. 17, 1992.

Field of the Invention

The present invention relates, in general, to a  
motility stimulating and compositions comprising the same.  
In particular, the present invention relates to a purified  
form of the protein and peptides thereof, for example,  
autotaxin (herein alternative referred to as "ATX"); a DNA  
segment encoding autotaxin; recombinant DNA molecules  
containing the DNA segment; cells containing the  
recombinant DNA molecule; a method of producing autotaxin;  
antibodies to autotaxin; and methods of cancer diagnosis  
and therapy using the above referenced protein or peptides  
thereof and DNA segments.

Background of the Invention

Cell motility plays an important role in  
embryonic events, adult tissue remodeling, wound healing,  
angiogenesis, immune defense, and metastasis of tumor  
cells (Singer, 1986). In normal physiologic processes,  
motility is tightly regulated. On the other hand, tumor  
cell motility may be aberrantly regulated or  
autoregulated. Tumor cells can respond in a motile  
fashion to a variety of agents. These include host-  
derived factors such as scatter factor (Rosen, et al.,  
1989) and growth factors (Kahan, et al., 1987; Stracke, et  
al.; Tamm, et al., 1989; Wang, et al. 1990; and Jouanneau,  
et al. 1991), components of the extracellular matrix  
(McCarthy, et al. 1984), and tumor-secreted or autocrine  
factors (Liotta, et al. 1988; Ruff, et al. 1985; Atnip, et  
al. 1987; Ohnishi, et al. 1990; Silletti, et al. 1991; and  
Watanabe, et al. 1991).

Many types of host-derived soluble factors act

- 2 -

in a paracrine fashion to stimulate cell locomotion. Motility-stimulating proteins called "scatter factors" have been identified which are produced by embryonic fibroblasts and by smooth muscle cells (Stoker, et al. 1987). Scatter factors stimulate random and directed motility by epithelial cells, keratinocytes, vascular endothelial cells and carcinoma cells (Stoker, et al. 1987; Rosen, et al. 1990; and Weidner, et al. 1990), but not fibroblasts. In addition, a number of host-secreted growth factors have been demonstrated to stimulate motility in tumor cells, including nerve growth factor (Kahan, et al. 1987) insulin-like growth factor-I (Stracke, et al. 1988), interleukin-6 (Tamm, et al. 1989), interleukin-8 (Wang, et al. 1990), and acidic fibroblast growth factor (Jouanneau, et al. 1991). These paracrine factors may influence "homing" or the directionality of tumor cell motility.

In contrast to these host-derived factors, many types of tumor cells have been found to produce proteins termed "autocrine motility factors" which stimulate motility by the same tumor cells which make the factor (Liotta, et al. 1986). Autocrine motility factors are not specific for a given type of cancer cell but have a wide spectrum of activity on many types of cancer cells (Kohn, et al. 1990), with little effect on normal fibroblasts or leukocytes.

Autocrine motility factors identified to date act through cell-surface receptors (Stracke, et al. 1987; Nabi, et al. 1990; Watanabe, et al. 1991) resulting in pseudopodial protrusion (Guirguis, et al. 1987) leading to both random and directed migration (Liotta, et al. 1986; Atnip, et al. 1987; Ohnishi, et al. 1990).

Prior studies of human A2058 melanoma cells have demonstrated that these cells are a particularly rich source of autocrine motility factors. An autocrine motility factor with a molecular mass of approximately 60



- 3 -

° kDa has been previously isolated from the conditioned media of these cells. (Liotta, et al. 1986). Similar tumor cells derived or induced factors with the same molecular weight have subsequently been reported and purified by several investigators (Atnip, et al. 1987; Schnor, et al. 1988; Ohnishi, et al. 1990; Silletti, et al. 1991; Watanabe et al. 1990). Such factors are thought to play a key role in tumor cell invasion.

Most of the motility factors identified to date have not been purified to homogeneity and have not been sequenced. The novel tumor motility factor of the present invention, named herein as autotaxin ("ATX"), has been purified and verified to be a homogeneous sample by two-dimensional gel electrophoresis. The protein of the present invention is unique from any previously identified or purified motility factor. The molecular size of ATX is about 125 kilo Daltons ("kDa") and it has an isoelectric point of approximately 7.7. ATX stimulates both random and directed migration of human A2058 melanoma cells at picomolar concentrations. The activity of the ATX factor is completely sensitive to inhibition by pertussis toxin. No significant homology has been found to exist between the protein of the invention and any mammalian protein including previous factors known to stimulate cell motility.

There is a great clinical need to predict the aggressiveness of a patient's individual tumor, to predict the local recurrence of treated tumors and to identify patients at high risk for development of invasive tumors. The present invention provides a functional marker which is functionally related to the invasive potential of human cancer. The invention further provides an assay for this secreted marker in body fluids, or in tissues. The assay of the invention can be used in the detection, diagnosis, and treatment of human malignancies and other inflammatory, fibrotic, infectious or healing disorders.

- 4 -

SUMMARY OF THE INVENTION

The present invention relates, generally, to a motility stimulating protein and corresponding peptides thereof, and to a DNA segment encoding same. A human cDNA clone encoding a tumor cell motility-stimulating protein, herein referred to as autotaxin or "ATX", reveals that this protein is an ecto/exoenzyme with significant homology to the plasma cell membrane differentiation antigen PC-1. ATX is a 125 kDa glycoprotein, previously isolated from a human melanoma cell line (A2058), which elicits chemotactic and chemokinetic responses at picomolar to nanomolar concentrations.

It is a specific object of the present invention to provide autotaxin and peptide fragments thereof.

It is a further object of the present invention to provide a DNA segment that encodes autotaxin and a recombinant DNA molecule comprising same. It is a further object of the present invention to provide a cell that contains such a recombinant molecule and a method of producing autotaxin using that cell.

Another object of the present invention is the identification of a transmembrane domain of the human liver autotaxin protein and its apparent absence in tumorous forms of autotaxin. The tumorous form of autotaxin appears to be a secreted protein. The present invention relates to utilization of the different sites of localization for diagnosis and prognosis of the stages of tumor progression. Further, the invention relates to treatment methods, designed to advantageously block the secreted form of autotaxin activity while having little effect on the membrane-bound form of autotaxin.

Yet another object of the present invention relates to the identification of a highly variable region within the autotaxin gene, called a "hot spot". The variations in sequence apparently result in mutations, insertions, deletions and premature termination of

- 5 -

translation. The present invention relates to manipulating this region so as to alter the activity of the protein. Further, the hot spot can serve as a marker in tumor diagnosis differentiating between different forms of the autotaxin protein.

It is yet another object of the present invention to provide a method of purifying autotaxin.

It is a further object of the present invention to provide cloned DNA segments encoding autotaxin and fragments thereof. The cDNA encoding the entire autotaxin protein contains 3251 base pairs, and has an mRNA size of approximately 3.3 kb. The full-length deduced amino acid sequence of autotaxin comprises a protein of 915 amino acids. Database analysis of the ATX sequence revealed a 45% amino acid identity (including 30 out of 33 cysteines) with PC-1, a pyrophosphatase/type I phosphodiesterase expressed on the surface of activated B cells and plasma cells. ATX, like PC-1, was found to hydrolyze the type I phosphodiesterase substrate p-nitrophenyl thymidine-5'-monophosphate. Autotaxin now defines a novel motility-regulating function for this class of ecto/exo-enzymes.

Further objects and advantages of the present invention will be clear from the description that follows.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Fractionation of ATX by hydrophobic interaction. A 200 ml sample of A2058 conditioned media was chromatographed on a 200 mL column of phenyl Sepharose-4B. Buffer A was 50 mM Tris (pH 7.5), 5% methanol, and 1.2 M ammonium sulfate. Buffer B was 50 mM Tris (pH 7.5), 5% methanol and 50% ethylene glycol. The gradient (----) represents a double linear gradient with decreasing ammonium sulfate (1.2 to 0.0 M) and increasing ethylene glycol (0 to 50%). Absorbance was monitored at 280 nm (—) and indicated that most of the protein did not bind to the column. Ten ml fractions were assayed for

- 6 -

motility stimulating capacity using the Boyden Chamber assay (o). The peak of motility activity occurred between 900 and 1050 minutes, ~ 12% of the gradient.

Figure 2. Isolation of ATX by lectin affinity chromatography. 20 ml portions of the phenyl Sepharose activity peak were affinity purified on a 40 ml Concanavalin A Affi-Gel column. The bound components were eluted with a step gradient (----) of methyl  $\alpha$ -D-mannopyranoside (0.0 mM, 10 mM, and 500 mM) in a buffer consisting of 0.05 M Tris (pH 7.5), 0.1 M NaCl, 0.01 M CaCl<sub>2</sub>, and 20% ethylene glycol. Absorbance was monitored at 280 nm (\_\_\_\_\_) and indicated that the majority of the protein components did not bind to the column. Motility was assayed in 10 mL fractions (...o...) and was found predominantly in the 500 mM elution concentration. One of seven chromatographic runs is shown.

Figure 3. Purification of ATX by weak anionic exchange chromatography. Approximately 30% of the activity peak eluted from the Con A affinity column was applied to a ZORBAX BioSeries-WAX column. The bound components were eluted with an NaCl gradient (----) in a buffer consisting of 10 mM Tris (pH 7.5) and 30% ethylene glycol. Motility (o) was assayed in 1.0 ml fractions. The peak of activity eluted in a discrete but broad region in the shallow portion of the gradient. Absorbance was monitored at 230 nm (\_\_\_\_). The majority of the protein components not associated with activity bound strongly to the column were eluted at 1.0 M NaCl. One of two chromatographic runs is shown.

Figure 4. Purification of ATX by molecular sieve exclusion chromatography. The entire activity peak eluted from the weak anion exchange column was applied to a series of TSK columns (4000SW, 4000SW, 3000SW, and 2000SW, in this order). Proteins were eluted in a buffer consisting of 0.1M NaPO<sub>4</sub> (pH 7.2) with 10% methanol and 10% ethylene glycol. Two major protein peaks were evident

- 7 -

by monitoring the absorbance at 235 nm (\_\_\_\_). Motility (...o...) was assayed in 0.4 ml samples and found predominantly in the first, smaller, protein peak.

Figure 5. Final purification of ATX by strong anionic exchange chromatography. Approximately 15% of the activity peak from the molecular sieve exclusion series was applied to a Pro-Pac PA1 column. Protein which bound to the column was eluted with a NaCl gradient (----) in a buffer consisting of 10 mM Tris (pH 7.5), 5% methanol and 20% ethylene glycol. Absorbance was monitored at 215 nm (\_\_\_\_). Motility activity was assayed in 1.0 ml fractions at two different dilutions: 1/5 (...o....) or 1/15 (.\_.o.\_.). Activity was found to correspond to a double protein peak in the central region of the gradient.

Figure 6A, 6B and 6C. Protein components associated with the activity peaks from various stages of purification. The activity peak from each chromatographic fractionation was pooled, concentrated and analyzed by SDS-polyacrylamide gel electrophoresis. Molecular weight standards are in Lane 1 for each panel. Panel 6A) 8-16% gradient gel of the first three purification steps, run under non-reducing conditions. Lane 2 is an aliquot of the pooled activity peak eluted from the phenyl sepharose fractionation. Lane 3 is an aliquot of the pooled activity peak eluted from the Con A affinity purification. Lanes 4 and 5 show the "peak" and "shoulder" of activity fractionated by weak anion exchange chromatography (Figure 3). Panel 6B) 7% gel of the activity peak fractionated by molecular sieve exclusion chromatography. Lanes 2 and 3 show the protein separation pattern of the total pooled activity peak when the gel was run under non-reducing and reducing conditions, respectively. Panel 6C) 8-16% gradient gel of the final strong anionic exchange chromatographic separation, run under non-reducing conditions. Lane 2 comprises ~1% of the total pooled activity peak eluted from the column.

- 8 -

Figure 7. Two-dimensional gel electrophoresis of ATX. Purified ATX (Figure 6, Panel C) was subjected to non-equilibrium isoelectric focusing (5 hr. at 500v), then applied to a 7.5% SDS-polyacrylamide gel for the second dimension. The pH separation which resulted was measured in 0.5 cm samples of concurrently run tube gels and is shown at the top. Molecular weight standards for the second dimension are shown on the right. This analysis reveals a single component with  $pI = 7.7 \pm 0.2$  and  $M_r = 120,000$ .

Figure 8. Dilution curve of ATX. Purified ATX (Figure 6, Panel C) was serially diluted and tested for motility-stimulating activity. The result, with unstimulated background motility subtracted out, shows that activity is half-maximal at  $\sim 500$  pM ATX.

Figure 9. Pertussis toxin (PT) sensitivity of ATX. A2058 cells were pre-treated for 1 hr. prior to the start of the motility assay with  $0.5 \mu\text{g/ml}$  PT in 0.1% BSA-DMEM or with 0.1% BSA-DMEM alone (for untreated control). The motility activity stimulated by purified ATX (Figure 6, Panel C) was then assessed for the two treatment groups. The result, expressed as cells/HPF  $\pm$  S.E.M. with unstimulated background motility subtracted out, reveals profound inhibition of PT-treated cells (hatched) compared to untreated cells (solid). PT had no effect on cell viability. S.E.M.'s were  $< 10\%$ .

Figure 10. Checkerboard analysis of ATX-stimulated motility. Varying dilutions of autotaxin were added to the upper chamber with the cells and/or to the lower chamber, as shown. Motility response, expressed as cells/HPF  $\pm$  S.E.M., was assessed for each point in the checkerboard.

Figure 11. Purification of ATX peptides on HPLC. ATX, purified to homogeneity by strong anionic exchange chromatography, was sequentially digested by cyanogen bromide, subjected to reduction and

- 9 -

pyridylethylation, and digested by trypsin. The resulting peptides were purified on an Aquapore RP300 C-8 reverse phase column using a (0-70)% acetonitrile gradient in 0.1% trifluoroacetic acid (----). The absorbance was monitored at 215 nm (\_\_\_\_) and peaks were collected. Seven peaks, chosen at random for N-terminal amino acid sequence analysis, are shown with appropriate numbers.

Figure 12. Cloning Strategy, schematically depicted.

Figure 13. Schematic Diagram of autotaxin gene region.

For A2058: 4C11 is the original DNA clone obtained by screening an A2058 cDNA expression library in  $\lambda$ gt11 with anti-peptide ATX-102. Upstream ATX peptide sequences were utilized for PCR amplification of A2058 mRNA, using the technique of reverse transcription/PCR. These peptides include ATX-101, ATX-103, and ATX-224. The approximate localization of each of peptide was obtained by matching the peptide with its homologous region on PC-1.

For N-tera 2D1, a  $\lambda$ gt10 cDNA library was amplified and the cDNA inserts were isolated. PCR amplification, based on homologies with A2058 sequence, was utilized for DNA sequencing.

For normal human liver, a mRNA from liver was amplified with 5'RACE using primers from the known ATX-224 region of A2058 and N-tera 2D1.

Figure 14. Schematic match-up of ATX peptides with putative A2058 protein sequence.

Figure 15. Schematic match-up of ATX peptides with putative N-tera 2D1 protein sequence.

Figure 16: ATX Treatment with PNGase F. Partially purified ATX was treated with 60 mU/ml PNGase F at 37°C for 16 hr under increasingly denaturing conditions. The treated ATX samples were separated by SDS polyacrylamide gel electrophoresis run under reducing

- 10 -

conditions and stained with Coomassie blue G-250. Lane 1 contains untreated ATX (arrow) with no enzyme added. Lane 2 contains the reaction mixture run under non-denaturing conditions (50 mM tris/10% ethylene glycol, pH 7). Lanes 3 and 4 have added 0.1 M  $\beta$ -mercaptoethanol and 0.5% Nonidet-P40, respectively. Lanes 5 and 6 contain the reaction mixtures in which the ATX sample was first boiled for 3 min in 0.1% SDS with (lane 6) or without (lane 5) 0.1 M  $\beta$ -mercaptoethanol, then had 0.5% Nonidet-P40 added to prevent enzyme denaturation. The enzyme can be detected as an ~44 kDa band in lanes 2-6.

Figure 17: Effect of varying concentrations of PNGase F on ATX molecular weight and motility-stimulating activity. Partially purified ATX was treated with various concentrations (range 0 - 60 mU/ml, shown on horizontal axis) of PNGase F at 37°C for 16 hr. Figure 17A shows the effect of the different treatments on ATX molecular weight. At concentrations of enzyme  $\geq 30$  mU/ml, the deglycosylation reaction appears to be complete. Figure 17B shows the effect of the identical reaction mixtures on motility-stimulating capacity (immediately below the corresponding protein band of Figure 17A). There is no significant difference between any of the treatment groups.

Figure 18: Comparison of amino acid sequences of ATX and PC-1. The amino acid sequences of ATX and PC-1 are compared. Amino acid identity is indicated by a vertical line between the sequences. The location of the putative transmembrane/signal sequence is shown by a solid line. The two somatomedin B domains are identified by dashed lines. The putative phosphodiesterase active site is indicated by emboldened lines. The loop region of a single EF hand loop region is identified with double lines. The presumed cleavage site for each protein is indicated with arrows.

Figure 19: Domain structure of ATX and PC-1.



- 11 -

Putative domains are indicated for the two homologous proteins, ATX and PC-1.

#### DETAILED DESCRIPTION OF THE INVENTION

Tumor cell motility is a critical component of invasion and metatasis, but the regulation of this motility is still poorly understood. At least some tumor cells secrete autocrine motility factors (AMF's) that stimulate motility in the producing cells. Like the analogous autocrine growth factors, these AMF's allow tumor cells independence from the host in this important component of the metastatic cascade. One AMF, autotaxin (ATX), has recently been purified to homogeneity from the human melanoma cell line, A2058 (Stracke, et al., 1992). The purified protein was enzymatically digested and the peptide fragments were separated by reverse phase HPLC. A number of these peptides have been sequenced by standard Edman degradation (Table 6) from different purifications and different enzymatic digestion. Sequence information, obtained initially on 19 purified tryptic peptides, confirmed that the protein is unique with no significant homology to growth factors or previously described motility factors. These peptide sequences have now been used as the basis for identifying and sequencing the cDNA clone for ATX. The present invention comprises an amino acid sequence of ATX as well as a nucleic acid sequence coding for the ATX protein.

TABLE 6. PEPTIDE SEQUENCES FOR AUTOTAXIN.

|    | PEPTIDE<br>NO. | AMINO ACID<br>SEQUENCE | SEQ ID:<br>NO: |
|----|----------------|------------------------|----------------|
| 30 | ATX-18         | WHVAR                  | SEQ ID NO:1    |
|    | ATX-19         | PLDVYK                 | SEQ ID NO:2    |
|    | ATX-20         | YPAFK                  | SEQ ID NO:3    |
|    | ATX-29         | PEEVTRPNYL             | SEQ ID NO:5    |
| 35 | ATX-34B        | RVWNYFQR               | SEQ ID NO:38   |

- 12 -

|    |              |                         |              |
|----|--------------|-------------------------|--------------|
| 0  | ATX-41       | HLLYGRPAVLY             | SEQ ID NO:29 |
|    | ATX-48       | VPPFENIELY              | SEQ ID NO:7  |
|    | ATX-59       | TFPNLYTFATGLY           | SEQ ID NO:32 |
|    | ATX-100      | GGQPLWITATK             | SEQ ID NO:8  |
| 5  | ATX-101/223A | VNSMQTVFVGYGPTFK        | SEQ ID NO:9  |
|    | ATX-102      | DIEHLTSLDFFR            | SEQ ID NO:10 |
|    | ATX-103      | TEFLSNYLTNVDDITLVPETLGR | SEQ ID NO:11 |
|    | ATX-104      | VNVISGPIDDYDYDGLHDTEDK  | SEQ ID NO:33 |
|    | ATX-204      | MHTARVRD                | SEQ ID NO:39 |
| 10 | ATX-205      | FSNNAKYD                | SEQ ID NO:40 |
|    | ATX-209      | VMPNIEK                 | SEQ ID NO:41 |
|    | ATX-210      | TARGWECT                | SEQ ID NO:42 |
|    | ATX-212      | (N)DSPWT(N)ISGS         | SEQ ID NO:43 |
|    | ATX-214      | LRSCGTHSPYM             | SEQ ID NO:44 |
| 15 | ATX-215/34A  | TYLHTYES                | SEQ ID NO:45 |
|    | ATX-213/217A | AIHANLTCKKPDQ           | SEQ ID NO:46 |
|    | ATX-216      | IVGQLMDG                | SEQ ID NO:47 |
|    | ATX-218/44   | TSRSYPEIL               | SEQ ID NO:48 |
| 20 | ATX-223B/24  | QAEVSSVPD               | SEQ ID NO:49 |
|    | ATX-224      | RCFELQEAGPPDDC          | SEQ ID NO:50 |
|    | ATX-229      | SYTSCCHDFDEL            | SEQ ID NO:51 |
|    | ATX-244/53   | QMSYGFLFPPYLSSSP        | SEQ ID NO:52 |

ATX is a glycosylated protein due to its high affinity for concanavalin A and amino acid sequence analysis of the ATX peptides. ATX has been demonstrated to be a 125kDa glycoprotein whose molecular weight reduced to 100-105kDa after deglycosylation with N-glycosidase F. The calculated molecular weight of the cloned protein is 100kDa (secreted form) or 105kDa (full length protein). Based on amino acid composition, the estimated pI is 9.0 which is higher than the pI determined by 2-D gel electrophoresis analysis (7.7-8.0) of purified ATX. This difference can be explained by the presence of sialic acid residues on the sugar moieties.

- 13 -

Autotaxin is secreted by A2058 human melanoma cells cultured in low abundance in serum-free conditioned medium. Autotaxin is a potent new cytokine with molecular mass 125 kDa which has been purified to homogeneity from the conditioned medium of the human melanoma cell line, A2058, utilizing sequential chromatographic methods as described herein. This new cytokine, termed autotaxin (ATX), is a basic glycoprotein with pI ~ 7.8. ATX is active in the high picomolar to low nanomolar range, stimulating both chemotactic and chemokinetic responses in the ATX-producing A2058 cells as well as other tumor cells. This motile response is abolished by pretreatment of the cells with pertussis toxin. ATX may therefore act through a G protein-linked cell surface receptor. These characteristics distinguish ATX from several small growth factors and interleukins which are implicated in tumor cell motility (Stracke et al., 1988; Ruff et al., 1985; Maciag et al., 1984; Gospodarowicz, 1984; Van Snick, 1990; Yoshimura 1987).

The protein of the present invention, which in one embodiment is derived from A2058 human melanoma cells, can be prepared substantially free from proteins with which it is normally associated using, for example, the purification protocol disclosed herein. Alternatively, the protein of the present invention can be prepared substantially free from proteins, by cloning and expressing the cDNA encoding autotaxin as disclosed herein.

A large volume of serum-free conditioned medium from appropriate producer cells (e.g., tumor cells) is collected and concentrated approximately 500-fold. This concentrated conditioned medium is then separated from other contaminating proteins by techniques that rely on the chemical and physical characteristics of the protein. These include the molecular weight, relative hydrophobicity, net charge, isoelectric focusing point,

- 14 -

° and the presence of lectin-binding sugar residues on the protein.

Alternatively, the protein, or functional portion thereof, can be synthesized using chemical or recombinant means.

5 The protein of the present invention has a potent biological activity. Purified ATX is active in the picomolar range and 1 unit of activity corresponds to a concentration of approximately 500 pM as assessed by the cell motility assay described herein and elsewhere  
10 (Stracke et al., 1989).

The protein of the present invention has a molecular size, as determined by two dimensional gel electrophoresis, of from 120 to 130 kDa, or more specifically, about 125 kDa. Further, the protein of the  
15 present invention can have a pI in the range of 7.5 to 8.0, preferably, approximately 7.7. The present invention relates to autotaxin and peptides thereof having cell motility properties as described herein. These proteins and peptides thereof can be produced by isolation from a  
20 natural host or isolation as an expression product from a recombinant host.

The present invention also relates to a DNA segment coding for a polypeptide comprising an amino acid sequence corresponding to ATX, or a unique portion of such a  
25 sequence (unique portion being defined herein as at least 5, 10, 25, or 50 amino acids). In one embodiment, the DNA segment encodes any one of the amino acid sequences shown in SEQ ID NO:1 to SEQ ID NO:11 and SEQ ID NO:26 to SEQ ID NO:33. Another embodiment uses larger DNA fragments  
30 encoding amino acid sequences shown in SEQ ID NO:34, SEQ ID NO: 36 and SEQ ID NO:38. The entire coding region for autotaxin can also be used in the present invention shown in SEQ ID NO:66 through SEQ ID NO:69.

35 In another embodiment, the present invention relates to a recombinant DNA molecule comprising a vector (for

- 15 -

° example plasmid or viral vector) and a DNA segment coding for a polypeptide corresponding to ATX, as can be prepared by one skilled in the art. Preferably, the coding segment is present in the vector operably linked to a promoter. The present invention also relates to a recombinant protein produced from a host cell expressing a cDNA  
5 containing a coding region of ATX. Examples of ATX cDNAs from a variety of sources have been cloned and can be used for expression, including *inter alia* A2058 carcinoma cells, N-tera 2D1 cells and human liver.

10 In a further embodiment, the present invention relates to a cell containing the above-described recombinant DNA molecule. Suitable host cells include procaryotic cells (such as bacteria, including *E. coli*) and both lower eucaryotic cells (for example, yeast) and  
15 higher eucaryotic cells (for example, mammalian cells). Introduction of the recombinant molecule into the host cells can be effected using methods known in the art.

In another embodiment, the present invention relates to a method of producing a polypeptide having an amino  
20 acid sequence corresponding to ATX. The method comprises culturing the above-described cells under conditions such that the DNA segment is expressed, and isolating ATX thereby produced.

In a further embodiment, the present invention  
25 relates to an antibody having affinity for ATX or peptide fragments thereof. The invention also relates to binding fragments of such antibodies. In one preferred embodiment, the antibodies are specific for ATX peptides having an amino acid sequence set forth in one of SEQ ID  
30 NO:1 through SEQ ID NO:11 and SEQ ID NO:26 through SEQ ID NO:34, SEQ ID NO: 36 and SEQ ID NO:38 through SEQ ID NO:52. In addition, the antibodies may recognize an entire autotaxin protein.

Antibodies can be raised to autotaxin or its fragment  
35 peptides, either naturally-occurring or recombinantly

- 16 -

produced, using methods known in the art.

The ATX protein and peptide fragments thereof described above can be joined to other materials, particularly polypeptides, as fused or covalently joined polypeptides to be used as carrier proteins. In particular, ATX fragments can be fused or covalently linked to a variety of carrier proteins, such as keyhole limpet hemocyanin, bovine serum albumin, tetanus toxoid, etc. See for example, Harper and Row, (1969); Landsteiner, (1962); and Williams et al., (1967), for descriptions of methods of preparing polyclonal antisera. A typical method involves hyperimmunization of an animal with an antigen. The blood of the animal is then collected shortly after the repeated immunizations and the gamma globulin is isolated.

In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts. Description of techniques for preparing such monoclonal antibodies may be found in Stites et al., and references cited therein, and in particular in Kohler and Milstein (1975), which discusses one method of generating monoclonal antibodies.

In another embodiment, the present invention relates to an oligonucleotide probe synthesized according to the sense or antisense degenerative sequence set forth in one of SEQ ID NO:1 through SEQ ID NO:11, SEQ ID NO:26 through SEQ ID NO:33, SEQ ID NO:39 through SEQ ID NO:52, and SEQ ID NO:55 through SEQ ID NO:65.

Protein database searches of this sequence revealed a 45% amino acid identity with the plasma cell membrane marker protein, PC-1. ATX and PC-1 appear to share a number of domains, including two somatomedin B domains, the loop region of an EF hand, and the enzymatic site of type I phosphodiesterase/ nucleotide pyrophosphatase. Like PC-1, ATX hydrolyzes p-nitrophenyl thymidine-5' - monophosphate, a type 1 phosphodiesterase substrate. This

- 17 -

- ° enzymatic function of ATX suggests a newly identified function for ecto/exo-enzymes in cellular motility.

In a further embodiment, the present invention relates to a method of diagnosing cancer metastasis and to a kit suitable for use in such a method. Preferably, antibodies to ATX can be used in, but not limited to, ELISA, RIA or immunoblots configurations to detect the presence of ATX in body fluids of patients (e.g. serum, urine, pleural effusions, etc.). These antibodies can also be used in immunostains of patient samples to detect the presence of ATX.

In yet another embodiment, the present invention relates to *in vivo* and *in vitro* diagnostics. ATX may be radiolabelled, by means known to one skilled in the art, and injected in cancer patients with appropriate ancillary substances also known to one skilled in the art, in order to ultimately detect distant metastatic sites by appropriate imagery. The level of ATX in tissue or body fluids can be used to predict disease outcomes and/or choice of therapy which may also include ATX inhibitors.

In a further embodiment, the present invention relates to a treatment of cancer. ATX antibodies can be cross-linked to toxins (e.g., Ricin A), by means known to one skilled in the art, wherein the cross-linked complex is administered to cancer patients with appropriate ancillary agents by means known to one skilled in the art, so that when the antibody complex binds to the cancer cell, the cell is killed by the cross-linked toxin.

In another embodiment, the different localizations of the normal versus tumorous forms of the ATX proteins within the tissue can be used as a tool for diagnosis and prognosis. The stage of disease progression can be monitored by elevated levels of ATX in the extracellular space as opposed to its normal cell membranes association. In addition, treatment methods for control of tumor progression can be designed to specifically block the

- 18 -

activity of the secreted form of ATX. Such methods would have a preferential effect upon secreted ATX during tumor progression while not effecting normal ATX formation.

Yet another embodiment utilizes the hot spot located in the region from approximately nucleotides 1670 through 1815, as a marker gene for identification of tissues carrying a tumorous form of ATX.

The present invention is described in further detail in the following non-limiting examples.

#### EXAMPLES

The following protocols and experimental details are referenced in the Examples that follow:

Materials. The polycarbonate Nuclepore membranes and the 48-well microchemotaxis chambers were obtained from Neuro Probe, Inc. Pertussis toxin (PT), ethylene glycol (biotechnology grade), methyl  $\alpha$ -D-mannopyranoside were obtained from commercial vendors. The ampholyte, pH 3-10 Bio-Lyte and pH 8-10 Bio-Lyte, were obtained from Bio-Rad. Phenyl Sepharose CL-4B; affi-Gel concanavalin A; ZORBAX BioSeries-WAX (weak anion exchange) column (9.4mm x 24cm); Spherogel-TSK 4000SW, 3000SW and 2000SW columns (each 7.5mm x 30cm); the Pro-Pac PA1 (4 x 50mm) strong anion exchange column; the Aquapore RP300 C-8 reverse phase column (220 x 2.1mm); and the AminoQuant C-18 reverse phase column (200 x 2.1mm) were also obtained from commercial sources.

Affi-Gel 10 affinity resin was from Bio-Rad. The GeneAmp PCR Reagent kit with AmpliTaq and the GeneAmp RNA PCR kit were purchased from Perkin-Elmer. The 5' RACE kit came from Gibco BRL Life Technologies, Inc. The p-nitrophenyl thymidine-5'-monophosphate was obtained from Calbiochem Biochemicals.

Ethylene glycol (biotechnology grade) was from Fisher Biochemicals (Pittsburg, PA). Peptide N-glycosidase F ("PNGase F"), O-glycosidase, neuraminidase (*Arthrobacter ureafaciens*), and swainsonine ("Swn") came from



- 19 -

Boehringer-Mannheim (Indianapolis, IN). 1-  
Deoxymannojirimycin ("dMAN"), and N-methyl-1-  
deoxynojirimycin ("NMdNM") were from Oxford GlycoSystems,  
Inc. (Rosedale, NY). Biotinylated concanavalin A, HRP-  
conjugated streptavidin, and HRP-conjugated goat anti-  
5 rabbit immunoglobulin were purchased from Pierce Chemicals  
(Rockford, IL). Polyvinyl pyrrolidone-free polycarbonate  
membranes and the microchemotaxis chamber were from  
NeuroProbe, Inc. (Cabin John, MD).

Cell Culture. The human melanoma cell line A2058,  
10 originally isolated by Todaro (Todaro et al., 1980), was  
maintained as previously described by Liotta (Liotta et  
al., 1986). The N-tera 2 (D1 clone) was a kind gift from  
Dr. Maxine Singer, Laboratory of Biochemistry, National  
Cancer Institute, National Institutes of Health and was  
15 maintained as described (Andrews, P.W., Goodfellow, P.N.  
and Bronson, D.L. (1983) *Cell surface characteristics and  
other markers of differentiation of human teratocarcinoma  
cells in culture.*).

Production of Autotaxin. A2058 cells were grown up in T-  
20 150 flasks, trypsinized, and seeded into 24,000 cm<sup>2</sup> cell  
factories at a cell density of  $1 \times 10^{10}$  cells/factory. After  
5-6 days, the serum-containing medium was removed and the  
cells were washed with DPBS. The factories were  
maintained in DMEM without phenol red, supplemented with 4  
25 mM glutamine, 100 units/ml penicillin, 100 µg/ml  
streptomycin, 5 µg/ml crystallized bovine serum albumin,  
10 µg/ml bovine insulin, and 1 µM aprotinin. Culture  
supernatants were harvested every 3 days, frozen at -40°C  
and replaced with fresh serum-free medium. Each cycle of  
30 supernatant was tested for ATX production with a cell  
motility assay detailed below. Typically, a cell factory  
continued to be productive for 9-11 of these cycles.

After accumulation of approximately 45-60 L of  
supernatant, the culture supernatants were thawed and  
35 concentrated down to 2-2.5 L using an Amicon S10Y30 spiral

- 20 -

° membrane ultrafiltration cartridge. This supernatant was further concentrated in an Amicon high performance ultrafiltration cell using Diaflo membranes. The final volume achieved from 100-200 L of conditioned medium was typically 250-400 ml. All ultrafiltrations were performed at 4°C.

5 Cell Motility Assays. Purification of autotaxin was monitored by testing the motility-stimulating capacity of the fractions collected from the columns. These fractions were in buffers unsuitable for a chemotaxis assay so each fraction had to be washed into an appropriate buffer, i.e., 0.1% (w/v) BSA in DPBS containing calcium and magnesium. This dialysis was performed by adding aliquots of each fraction to be tested into Centricon-30™ ultrafiltration tubes, which retain molecular species larger than 30,000 daltons.

15 The assay to determine motility was performed in triplicate using a 48-well microchemotaxis chamber as described elsewhere in detail (Stracke et al., 1987; Stracke, et al., 1989). The Nuclepore™ membranes used in these modified Boyden chambers were fixed and stained with Diff-Quik.™ Chemotaxis was quantitated either by reading the stained membranes with a 2202 Ultrosan laser densitometer or by counting 5 randomly chosen high power fields (HPF) under light microscopy (400 x) for each replicate. Densitometer units (wavelength - 633 nm) have been shown to be linearly related to the number of cells per HPF (Taraboletti, 1987; Stracke, et al., 1989). Typically, unstimulated motility (background) corresponded to 5-10 cells/HPF and highly responding cells to 70-100 cells/HPF above unstimulated background (i.e., 75-110 total cells/HPF).

20 For experiments using PT, the toxin was pre-incubated with the cells for 1-2 hr. at room temperature prior to the assay and maintained with the cells throughout the assay (Stracke, et al., 1987). The treated cells were

25

30

35

- 21 -

- ° tested for their motility response to the chemoattractant as well as for unstimulated random motility.

Purification of Autotaxin. Ammonium sulfate, to a final concentration of 1.2 M, was added to the concentrated A2058 conditioned medium for 1 hr. at 4°C. The solution  
5 was spun in a RC2-B Ultraspeed Sorvall centrifuge at 10,000 x g for 15 min. Only the supernatant had the capacity to stimulate motility.

In the first step, the sample was fractionated by hydrophobic interaction chromatography using 200 ml phenyl  
10 Sepharose CL-4B column equilibrated into 50 mM Tris (pH 7.5), 5% (v/v) methanol and 1.2 M ammonium sulfate. The supernatant from the ammonium sulfate fractionation was added to this column and eluted using linear gradients of 50 mM Tris (pH 7.5), 5% (v/v) methanol, with decreasing  
15 (1.2 - 0.0) M ammonium sulfate and increasing (0-50) % (v/v) ethylene glycol at 1 ml/min.

The active peak was pooled, dialyzed into 50 mM Tris, 0.1 M NaCl, 0.01 M CaCl<sub>2</sub>, 20% (v/v) ethylene glycol, and subjected to a second fractionation by lectin affinity  
20 chromatography using a 40 ml Affi-Gel concanavalin A column run at 1 ml/min. The sample was eluted in a stepwise fashion in the same buffer with 0, 10, and 500 mM methyl  $\alpha$ -mannopyranoside added successively. Fractions from each step of the gradient were pooled and tested for  
25 their capacity to stimulate motility.

In the third purification step, the sample that eluted at 500 mM  $\alpha$ -methyl-mannopyranoside was dialyzed into 10 mM Tris (pH 7.5) with 30% (v/v) ethylene glycol and fractionated by weak anion exchange chromatography.  
30 Chromatography was carried out on a ZORBAX BioSeries-WAX column using a Shimadzu BioLiquid chromatograph and eluted with a linear gradient of (0.0 - 0.4 M) sodium chloride at 3 ml/min.

The active peak was pooled, dialyzed against 0.1 M  
35 sodium phosphate (pH 7.2), 10% (v/v) methanol, and 10%

- 22 -

° (v/v) ethylene glycol, and subjected to a fourth fractionation step on a series of Spherigel TSK columns (4000SW, 4000SW, 3000SW, 2000SW, in that order). This molecular sieve step was run using the Shimadzu BioLiquid chromatograph at 0.4 ml/min.

5 The active peak was pooled and dialyzed into 10 mM Tris (pH 7.5), 5% (v/v) methanol, 20% (v/v) ethylene glycol and subjected to a fifth (strong anion exchange) chromatography step, a Pro-Pac PA1 column run at 1 ml/min using a Dionex BioLC with AI450 software. The sample was  
10 eluted with a linear gradient of (0.0-0.4M) NaCl.

In order to calculate activity yields after each step of purification, a unit of activity had to be derived. The dilution curve of ATX was biphasic with a broad peak and a linear range at sub-optimal concentrations. One unit  
15 of activity/well (i.e., 40 units/ml) was defined as 50% of the maximal activity in a full dilution curve. This allowed calculation of the activity contained in any volume from the dilution needed to achieve 1 unit/well. Therefore, if a 1:10 dilution were needed in order to  
20 produce 1 unit of activity/well, the material contained  $10 \times 40 = 400$  units/ml.

Gel Electrophoresis. Protein samples were analyzed by SDS-polyacrylamide gel electrophoresis using the conditions of Laemmli (Laemmli, 1970). In brief, 7 or 8% SDS-containing  
25 polyacrylamide gels were prepared or pre-poured (8-16%) gradient gels were obtained commercially. Samples were prepared with or without reducing conditions (5%  $\beta$ -mercaptoethanol). After electrophoretic separation, the gels were stained using Coomassie Blue G-250 as previously  
30 described (Neuhoff, et al., 1988). In this staining protocol, which ordinarily requires no destaining step, the Coomassie stain appears to be able to stain as little as 10 ng of protein.

For two-dimensional electrophoresis, the protein, in  
35 20% ethylene glycol, was dried in a Speed-vac and

- 23 -

redissolved in loading solution: 9M urea, 1% (v/v) pH 3-10 Bio-Lyte, and 2.5% (v/v) Nonidet-P40. This sample was then subjected to isoelectric focusing (O'Farrell, 1975) using a Bio-Rad tube cell in 120 x 3 mm polyacrylamide tube gels containing 9M urea, 2% (v/v) pH 3-10 Bio-Lyte, 0.25% (v/v) pH 8-10 Bio-Lyte and 2.5% (v/v) Nonidet-P40. Reservoir solutions were 0.01 M phosphoric acid and 0.02 M NaOH. Non-equilibrium isoelectric focusing (O'Farrell, et al., 1977) was run initially with constant voltage (500 v) for 5 hr. Since the protein was basic, the procedure was repeated under equilibrium conditions (500 v for 17 hr.). Electrophoresis in the second dimension was performed on a 7.5% polyacrylamide using the conditions of Laemmli (1970). The gel was stained with Coomassie Blue G-250 as above.

Preparation of peptides for internal sequence of autotaxin. Homogeneous ATX was sequentially digested with cyanogen bromide and, following reduction and pyridylethylation, with trypsin (Stone, et al., 1989). The resulting fragments were then separated by gradient elution on an Aquapore RP300 C-8 reverse phase column: 0.1% (v/v) trifluoroacetic acid and (0-70)% acetonitrile over 85 min. at a flow rate of 0.2 ml/min. A Dionex AI450 BioLC system was utilized and fractions were collected manually while monitoring the absorbance at 215 nm.

Sequence analysis of peptides. The amino acid sequences of peptides resulting from digestion and purification of ATX peptides #1-7 and 12-18, corresponding to SEQ ID NO:1 through SEQ ID NO:7 and SEQ ID NO:26 through SEQ ID NO:32, respectively, were determined on a Porton Instruments 2020 off-line sequenator using standard program #1. Phenylthiohydantoin amino acid analysis of sequenator runs were performed on a Beckman System Gold HPLC using a modified sodium acetate gradient program and a Hewlett-Packard C-18 column. ATX-100 (SEQ ID NO:8), ATX-101 (SEQ ID NO:9), ATX-102 (SEQ ID NO:10), ATX-103 (SEQ ID NO:11)

- 24 -

- and ATX 104 (SEQ ID NO:33) were sequenced from gel-purified ATX.

Protein databases (Pearson, et al. 1988) that were searched for homologies in amino acid sequence with the ATX peptides include: GenBank (68.0), EMBL (27.0), SWISS-PROT (18.0), and GenPept (64.3).

#### EXAMPLE 1

##### Purification of Autotaxin

The A2058 cells had been previously shown to produce protein factors which stimulate motility in an autocrine fashion (Liotta, et al., 1986). Conditioned medium from these cells was therefore used to identify and purify a new motility-stimulating factor, which is here named autotaxin and referred to as ATX. Since the purification was monitored with a biological assay, motility-stimulating activity had to be maintained throughout. The activity proved to be labile to freezing, acidic buffers, proteases (but not DNase or RNase), reduction, strong chaotropic agents (e.g. > 4 M urea), and a variety of organic solvents (isopropanol, ethanol, acetonitrile). An organic solvent, ethylene glycol, which did not decrease bioactivity, was added for both storage and chromatographic separation.

100-200 L of serum-free conditioned medium were required in order to produce enough ATX for amino acid sequence analysis. The medium contained low concentrations of both BSA (5  $\mu$ g/ml) which was needed as a carrier protein and insulin (10  $\mu$ g/ml) which was required to support cell growth in low protein medium. Ultrafiltration to concentrate this large volume was performed with low protein-binding YM30 membranes which retain molecular species with  $M_r$  > 30,000. As seen in Table 1, 200 L of conditioned medium prepared in this manner resulted in  $10 \times 10^6$  units of activity. However, the initial unfractionated conditioned medium contained additional substances known to stimulate activity,

- 25 -

- ° particularly insulin, which does not completely wash out in the ultrafiltration step and which is additive to the motility stimulating activity in a complex manner (Stracke, et al., 1989). This had to be taken into account in order to determine yields for subsequent steps in which insulin had been removed.

TABLE 1. PURIFICATION OF AUTOTAXIN

| 10 | Purification Step<br>Recovery | Protein<br>(mg)    | Activity <sup>a</sup><br>(total units) | Specific               |                  |
|----|-------------------------------|--------------------|--|------------------------|------------------|
|    |                               |                    |  | Activity<br>(units/mg) | (%) <sup>b</sup> |
|    | 200 L Conditioned Medium      | 33,000             | 10,000,000 <sup>c</sup>                | 300                    |                  |
|    | Phenyl Sepharose              | 1,235              | 460,000                                | 370                    | 100              |
| 15 | Concanavalin A                | 58                 | 660,000                                | 11,400                 | 100              |
|    | Weak Anion Exchange           | 4.5                | 490,000                                | 110,000                | 100              |
|    | TSK Molecular Sieves          | ~0.4 <sup>d</sup>  | 220,000                                | 550,000                | 48               |
| 20 | Strong Anion Exchange         | ~0.04 <sup>d</sup> | 24,000 <sup>e</sup>                    | 600,000                | 5.2              |

<sup>a</sup> Activity calculated from Boyden chamber assay. The dilution which resulted in 50% of maximal activity (generally approximately 20 laser density units or ~40 cells/HPF) was chosen to have 1 unit of activity per well (equivalent to 40 units/ml).

25 <sup>b</sup> Recovery was estimated from activity, after the first purification column (i.e., phenyl sepharose).

<sup>c</sup> Initial activity in the unfractionated conditioned medium reflected the fact that insulin was used in the medium as a necessary growth factor under low protein conditions.

<sup>d</sup> Estimated protein is based on quantification by amino acid analysis.

30 <sup>e</sup> This specific activity for purified protein corresponds to ~10 fmol ATX/unit of motility activity (in a Boyden chamber well).

The first step in the purification involved fractionation by hydrophobic interaction chromatography using a phenyl Sepharose CL-4B column. The results are

- 26 -

shown in Figure 1. Most proteins, including insulin, eluted from the column in early fractions or in the void. However, the peak of activity eluted relatively late. The activity which was purified was estimated as 460,000 units  $\pm$  20% (Table 1). As the pooled peak of activity from the phenyl Sepharose fractionation is considered to be the first sample without significant insulin contamination, subsequent yields are measured against its total activity. Gel electrophoresis of a small portion of the pooled peak of activity (Figure 6A, column 2) revealed a large number of protein bands with BSA predominant from the original conditioned medium.

In the second step of purification, the active peak was applied to the lectin affinity column, Affi-Gel concanavalin A. As shown in Figure 2, most protein (estimated to be 90% of the total absorbance at 280 nm) failed to bind to the column at all. The non-binding fraction contained essentially no motility-stimulating activity (see dotted line in Figure 2). When a linear gradient of methyl  $\alpha$ -D-mannopyranoside was applied to the column, chemotactic activity eluted off in a prolonged zone, beginning at a concentration of approximately 20 mM sugar. Consequently, a step gradient was used to elute. Pure BSA failed to bind to con A.

Activity was found primarily in the 500 mM step of methyl  $\alpha$ -D-mannopyranoside. There appeared to be no significant loss of activity as seen in Table 1; however, specific activity (activity/mg total protein) increased thirty-fold. Gel electrophoresis of the pooled and concentrated peak (Figure 6A, column 3) revealed that the BSA overload was no longer apparent and the number of bands were much reduced. When the unbound protein was concentrated and applied to a gel, it appeared identical to the active peak from phenyl Sepharose-4B with a large BSA band.

The third purification step involved



- 27 -

fractionating the previous active peak by weak anion exchange chromatography as shown in Figure 3. Under the running conditions, activity eluted in a broad peak-shoulder or double peak in the middle of the shallow portion (0.0-0.4 M) of the NaCl gradient. The largest proportion of protein, lacking in motility-stimulating capacity, bound strongly to the column and eluted off in high salt (1 M NaCl). There appeared to be no significant loss of activity, though specific activity increased by twenty-fold (Table 1). Analysis by gel electrophoresis of both the peak (28-34 min. in Figure 3) and the shoulder (35-45 min. in Figure 3) is shown in Figure 6A (columns 4 and 5, respectively). Two predominant protein bands resulted: a broad doublet around 25-35 kDa and a second doublet around 110-130 kDa.

In the fourth purification step, the active peak was applied to a series of molecular sieves. Spectrophotometric monitoring of the eluant revealed two large peaks of protein (Figure 4). Activity corresponded to the first, higher molecular weight peak. Recovery of activity was ~48% with a five-fold increase in specific activity. Analysis by gel electrophoresis was performed under non-reducing and reducing conditions as shown in Figure 6B (columns 2 and 3, respectively). This fractionation step had essentially removed all contaminating protein of molecular weight < 55 kDa. The predominant band remaining has a molecular weight of 120 kDa unreduced and 125 kDa reduced; there are two minor bands with molecular weights 85 kDa and 60 kDa. The fact that the 120 kDa protein changes so little in electrophoretic mobility after reduction would tend to indicate a paucity of disulfide bonds. However, the existing disulfide bonds have functional significance because motility-stimulating activity is labile to reduction.

The fifth purification step involved

- 28 -

fractionation of the active peak by strong anion exchange chromatography. As shown in Figure 5, activity corresponds to two broad optical absorbance peaks in the middle of the gradient with contaminating proteins eluting earlier. These two peaks were identical by amino acid analysis and by polyacrylamide gel electrophoretic separation. They presumably represent different glycosylation states of the same parent protein. Activity is shown in Figure 5 at two different sample dilutions. Several dilutions of the fractionated samples were often necessary in order to resolve the true "peak" of activity as the shape of the ATX dilution curve was not sharp due to saturation and down regulation at high concentrations. Recovery from this chromatographic step is lower (5% compared to phenyl Sepharose), as might be expected when a minute quantity of protein is applied to a column; however, specific activity again increased (Table 1). Analysis by gel electrophoresis revealed a single protein band at molecular weight 120 kDa, unreduced (Figure 6C, column 2).

#### EXAMPLE 2

##### Characterization of Autotaxin

Two dimensional gel electrophoresis of the purified protein (Figure 7) revealed a single predominant band. The band slopes downward slightly toward the basic side of the gel in a manner that is characteristic of glycosylated proteins. A basic pI of  $7.7 \pm 0.2$  was essentially the same whether the isoelectric focusing was run under non-equilibrium conditions (5 hr.) or was allowed to go to equilibrium (17 hr.).

A dilution curve of the purified protein is shown in Figure 8. The protein is active in the picomolar range and 1 unit of activity appears to correspond to a concentration of 400-600 picomolar (or approximately 10 fmol of ATX/Boyden chamber well). When dilutions were begun at higher concentrations of ATX, the resultant curve showed a broad plateau with down-regulation at the highest

- 29 -

concentrations. The motility response to purified autotaxin is highly sensitive to pertussis toxin (hereinafter referred to as "PT") (Table 2 and Figure 9) with approximately 95% inhibition of activity at 0.5  $\mu\text{g/ml}$  PT.

TABLE 2. Effect of Pertussis Toxin (PT) on Autotaxin-stimulated motility

|                               | A2058 Motility Response<br>(density units <sup>1</sup> ) |  |
|-------------------------------|--|--|
|                               | control cells <sup>2</sup>                               | Pertussis toxin-treated cells <sup>3</sup> |
| Condition medium <sup>4</sup> | 60.3   | 0.4  |
| Purified Autotaxin            | 38.5   | 0.0  |

<sup>1</sup> Chemotaxis quantitated by motility assay (Stracke, et al., 1978).

<sup>2</sup> A2058 cell suspended at  $2 \times 10^6$  cells/ml in DMEM supplemented with 1 mg/ml bovine serum and rocked at room temperature for 1 hr.

<sup>3</sup> As control with 0.5  $\mu\text{g/ml}$  pertussis toxin.

<sup>4</sup> Prepared by adding DMEM without phenol red supplemented with 0.1 mg/ml bovine serum albumin to subconfluent flasks of A2058 cells. The medium was harvested after 2 days incubation at 37°C in a humidified atmosphere and concentrated 25-30 fold using an Amicon ultrafiltration assembly with a YM-30 membrane.

Checkerboard analysis was performed to assess the random (chemokinetic) versus the directed (chemotactic) nature of the motility response to ATX. Chambers were assembled with different concentrations of ATX above and below the filter, using ATX purified through the weak anion exchange fractionation step. Squares below the diagonal reflect response to a positive gradient, squares above reflect response to a negative gradient, and squares on the diagonal reflect random motility in the absence of a gradient. ATX stimulates both chemotactic and chemokinetic responses (Figure 10), with chemotactic responses as high as fifteen-fold above background and

- 30 -

chemokinesis as high as eight-fold above background.

Amino acid analysis after complete acid hydrolysis was used to quantitate purified protein. This hydrolysis was carried out on protein excised from a polyacrylamide gel and presumed to be pure. The analysis indicated that 2.7 nmol of protein was present after fractionation on the molecular sieve. After fractionation by strong anion exchange chromatography, approximately 300 pmol remained. The results of the analysis are shown in Table 3.

TABLE 3. AMINO ACID COMPOSITION OF AUTOTAXIN  
(CYS and TRP were not determined in this analysis)

| <u>Amino Acid</u> | <u>Residues/100</u> |
|-------------------|---------------------|
| ASX               | 12.5                |
| THR               | 6.0                 |
| SER               | 5.7                 |
| GLX               | 9.4                 |
| PRO               | 7.4                 |
| GLY               | 7.0                 |
| ALA               | 3.9                 |
| VAL               | 6.7                 |
| MET               | 1.2                 |
| ILE               | 4.3                 |
| LEU               | 9.0                 |
| TYR               | 5.2                 |
| PHE               | 5.2                 |
| HIS               | 3.8                 |
| LYS               | 7.4                 |
| ARG               | 5.4                 |

### EXAMPLE 3

#### ATX Degradation and Determination of Amino Acid Sequence

Attempts to obtain N-terminal sequence information from purified ATX repeatedly proved futile. The purified protein was therefore, sequentially digested and the resulting peptides fractionated by reverse phase chromatography. The results are shown in Figure 11. Multiple sharp peaks including clusters at both the hydrophilic and hydrophobic ends of the gradient are seen.

- 31 -

Several of these peptide peaks were chosen randomly for Edman degradation and N-terminal amino acid sequence analysis. Seven of the eight peaks (shown in Figure 11) chosen gave clear single sequence information as seen in Table 4. Using material from a separate digestion and purification, the remaining four sequences were also obtained.

Separate sense and antisense oligonucleotide probes were synthesized according to the fragment sequences of Table 4 by methods known to one skilled in the art. Representative probes are shown in Table 5.

TABLE 4. Peptide sequences for Autotaxin.

|    | PEPTIDE<br>NO. | AMINO ACID<br>SEQUENCE       | SEQ ID:<br>NO: | NAME    |
|----|----------------|------------------------------|----------------|---------|
| 15 | 1.             | WHVA                         | SEQ ID NO:1    | ATX 18  |
|    | 2.             | PLDVYK                       | SEQ ID NO:2    | ATX 19  |
|    | 3.             | YPAFK                        | SEQ ID NO:3    | ATX 20  |
|    | 4.             | QAEVS                        | SEQ ID NO:4    | ATX 24  |
|    | 5.             | PEEVTRPNYL                   | SEQ ID NO:5    | ATX 29  |
| 20 | 6.             | YDVPWNETI                    | SEQ ID NO:6    | ATX 47  |
|    | 7.             | VPPFENIELY                   | SEQ ID NO:7    | ATX 48  |
|    | 8.             | GGQPLWITATK                  | SEQ ID NO:8    | ATX 100 |
|    | 9.             | VNSMQTVFVGY-<br>GPTFK        | SEQ ID NO:9    | ATX 101 |
| 25 | 10.            | DIEHLTSLDFFR                 | SEQ ID NO:10   | ATX 102 |
|    | 11.            | TEFLSNYLTNVDD-<br>ITLVPETLGR | SEQ ID NO:11   | ATX 103 |
|    | 12.            | QYLHQYGSS                    | SEQ ID NO:26   | ATX 37  |
| 30 | 13.            | VLNYF                        | SEQ ID NO:27   | ATX 39  |
|    | 14.            | YLNAT                        | SEQ ID NO:28   | ATX 40  |
|    | 15.            | HLLYGRPAVLY                  | SEQ ID NO:29   | ATX 41  |
|    | 16.            | SYPEILTPADN                  | SEQ ID NO:30   | ATX 44  |
|    | 17.            | XYGFLFPPYLSSSP               | SEQ ID NO:31   | ATX 53  |
| 35 | 18.            | TFPNLYTFATGLY                | SEQ ID NO:32   | ATX 59  |

- 32 -

19. VNVISGPIFDYDYDGLH SEQ ID NO:33 ATX 104  
DTEDK

Peptide numbers 1-7 refer to peaks numbered in Figure 11. Peptide numbers 12-18 refer to peptides purified from the preparation which yielded peptide numbers 1-7. Peptides 8-11 and 19, are from a separate purification, not shown in Figure 11.

X refers to potentially glycosylated residues.

TABLE 5.

10 Oligonucleotides synthesized from peptide sequences of autotaxin (ATX). The number of the oligonucleotide corresponds to the ATX peptide number as per Table 4. The final letter suffix distinguishes whether the oligonucleotide is a sense (S) or antisense (A) sequence.

|    | <u>Oligo</u> | <u>Sequence</u>   | <u>SEQ ID NO:</u> |
|----|--------------|---|-------------------|
| 15 | A-18A        | GTT-GGC-AGC-NAC-RTG-CCA                                 | SEQ ID NO:12      |
|    | A-18S        | TGG-CAY-GTN-GCT-GCC-AAC                                 | SEQ ID NO:13      |
|    | A-20A        | CTT-GAA-GGC-AGG-GTA                                     | SEQ ID NO:14      |
|    | A-20S        | TAY-CCT-GCN-TTY-AAG                                     | SEQ ID NO:15      |
|    | A-29A        | GGT-NAC-YTC-YTC-AGG                                     | SEQ ID NO:16      |
| 20 | A-29S        | CCT-GAR-GAR-GTN-ACC                                     | SEQ ID NO:17      |
|    | A-47A        | NGT-NGC-RTC-RAA-TGG-CAC-RTC                             | SEQ ID NO:18      |
|    | A-47S        | GAY-GTG-CCA-TTY-GAY-GCN-ACN                             | SEQ ID NO:19      |
|    | A-48A        | GTT-DAT-RTT-STC-RAA-TGG-GGG                             | SEQ ID NO:20      |
|    | A-48S        | CCC-CCA-TTT-GAG-AAC-ATC-AAC                             | SEQ ID NO:21      |
| 25 | A-100A       | CTT-NGT-NGC-NGT-DAT-CCA-NAR-<br>GGG-YTG-GCC-GCC         | SEQ ID NO:22      |
|    | A-100S       | GGC-GGC-CAR-CCC-YTN-TGG-ATH-<br>ACN-GCN-ACN-AAG         | SEQ ID NO:23      |
| 30 | A-101A       | CTT-RAA-GGT-GGG-GCC-RTA-GCC-<br>CAC-RAA-GAC-TGT-YTG-CAT | SEQ ID NO:24      |
|    | A-101S       | ATG-CAR-ACA-GTC-TTY-GTG-GGC-<br>TAY-GGC-CCC-ACC-TTY-AAR | SEQ ID NO:25      |

35

- 33 -

EXAMPLE 4Antipeptide Antibodies

Rabbits were injected with ATX-101 (SEQ ID NO:10) which had been cross-linked to bovine serum albumin. Antisera from these rabbits was subjected to salt precipitation followed by purification using affinity chromatography with Affi-Gel 10 beads covalently linked to the peptide, ATX-101 (SEQ ID NO:10). This affinity purified antibody reacted with the partially purified protein on immunoblots. This same antibody has been used to perform immunohistochemical stains on human tissue.

EXAMPLE 5Enzymatic Deglycosylation of ATX

Purified ATX that was to be treated with peptide N-glycosidase F (PNGase F) was first dialyzed into 0.2 M sodium phosphate, 10% (v/v) ethylene glycol pH 7.0, using Centricon-30 ultrafiltration tubes. Varying concentrations of PNGase F were added to the ATX and incubated 16-18 hr. at 37°C. Complete digestion appeared to occur at concentrations of enzyme above 30 mU/ml (where 1 U converts 1mmol of substrate/min). For comparison, the experiments were repeated in the presence of 0.1 M  $\beta$ -mercaptoethanol or 0.1% (w/v) SDS plus 0.5% (v/v) Nonidet-P40. ATX that was to be treated with neuraminidase or O-glycosidase was dialyzed into 20 mM sodium phosphate, 0.1 M calcium acetate, and 10% (v/v) ethylene glycol (pH 7.2). Neuraminidase was added to a final concentration of 2 U/ml. For treatment with neuraminidase alone, this mixture was incubated 16-18 hr at 37°C. Since O-glycosidase requires the removal of terminal sialic acid residues for efficient deglycosylation, ATX was pre-incubated with neuraminidase for 30-125 mU/ml and incubated 16-18 hr. at 37°C. The treated ATX was then dialyzed into 50 mM Tris with 20% ethylene glycol for storage at 5°C.

35

- 34 -

°     Treatment of ATX with N-glycosylation altering agents

A2058 cells were split into four 150 cm<sup>2</sup> flasks and incubated until just subconfluent in DMEM supplemented with 10% fetal calf serum. The medium was then replaced with fresh 10% FCS/DMEM to which had been added DPBS for control, 1mM dMAN, 1 mM NMdNM, or 10 mM (1.7 mg/ml) Sw<sub>n</sub>. Concentrations of these pharmacological agents were similar to those previously described as inhibiting N-glycan processing enzymes in melanoma cells (Seftor, et al. 1991; Dennis, et al. 1990) as well as carcinoma cells (Ogier, et al. 1990). On the next day, each flask was washed twice with Dulbecco's phosphate buffered saline with calcium ("DPBS") then 20 ml of Dulbecco's minimum essential medium ("DMEM") supplemented with 0.01% (w/v) bovine serum albumin ("BSA") was added. The same concentration of each agent was added to the appropriate equilibrated flask and incubated for ~ 24 hr, after which the medium from each treatment group was collected, concentrated, washed into DPBS and stored at 5°C.

Cells from each flask were trypsinized and counted. There was no loss of viability or reduced cell number in any of the treatment groups compared to control cells.

Effect of PNGase F on ATX

ATX binds to concanavalin A ("Con A") agarose beads and is eluted with buffer containing 0.5 M methyl α-D-mannopyranoside, indicating that ATX is likely to contain mannose residues. Such mannose sugar residues are most characteristic of N-linked oligosaccharides. In order to verify that ATX contained asparagine-linked oligosaccharides, we treated it with the endoglycosidase, PNGase F, which cleaves high mannose, hybrid, and complex N-linked oligosaccharides at the asparagine residue.

Partially purified ATX was treated with 60 mU/ml of enzyme under a variety of increasingly denaturing conditions and then separated by polyacrylamide gel



- 35 -

electrophoresis (Figure 16). Lane 1 shows untreated material; the 125 kDa band (arrow) is autotaxin. When this material is treated overnight with PNGase F under very mild conditions, the size of the 125 kDa band decreases to ~100- 105 kDa. Addition of 0.1 M b-mercaptoethanol (Lane 2) or 0.5% Nonidet-P40 (lane 3) to the ATX sample has no effect on the size of the resultant protein band. Even complete denaturation of ATX of boiling the sample for 3 min in 0.1% SDS with (lane 5) or without (lane 4)  $\beta$ -mercaptoethanol, followed by addition of 0.5% Nonidet-P40 to maintain enzymatic activity, has no effect on the final size of deglycosylated protein, indicating that the deglycosylation reaction was complete even under mild conditions.

Because these results showed that ATX contained N-linked oligosaccharide groups, it became important to see if these sugar moieties were necessary for stimulation of motility. The partially purified ATX sample was treated with varying concentrations of PNGase F (0.1 to 60 mU/ml) under mild, non-denaturing conditions. Analysis of the resulting digest by polyacrylamide gel electrophoresis is shown in Figure 17A. As this figure shows, the digestion was incomplete using from 0.1 to 10 mU/ml of enzyme and resulted in a smear of protein between 100-125 kDa. However, at higher concentrations of enzyme, cleavage of N-linked oligosaccharides from ATX appears to be complete. When these different digestion products were compared for their capacity to stimulate motility (Figure 17B), there was no significant difference between groups.

#### EXAMPLE 6

##### Cloning the 3' end of Autotaxin (4C11)

ATX is active in picomolar to nanomolar concentrations and is synthesized in very small concentrations by A2058 cells. As might be expected, the cDNA clone was relatively rare, requiring various strategies and multiple library screenings in order to

- 36 -

° identify it (Figure 12). Attempts to utilize degenerate oligonucleotides deduced from known peptide sequences were unsuccessful--whether we used the oligo nucleotides for screening cDNA libraries or for reverse transcription of mRNA followed by amplification with the polymerase chain reaction (RT/PCR). We then utilized an affinity-purified anti-peptide ATX-102 antibodies to screen an A2058 expression library.

These anti-peptide antibodies were generated by methods well established in the art and described previously with slight modification (Wacher, et al., 1990). In brief, the previously identified peptide, ATX-102 (Stracke, et al., 1992), was synthesized on a Biosearch 9600 peptide synthesizer. It was then solubilized in 1X PBS containing 20% (v/v) DMSO and conjugated to the protein carrier, bovine serum albumin (BSA), with glutaraldehyde. For the first injection into New Zealand white rabbits, the BSA-peptide conjugate was emulsified with complete Freund's adjuvant and injected subcutaneously. For subsequent injections, the BSA-peptide conjugate was emulsified with incomplete Freund's adjuvant. The resultant antiserum was heat-inactivated at 56°C for 30 min. Immunoglobulins were precipitated out in 47% saturated ammonium sulfate, then redissolved and dialyzed into PBS. Antibodies were adsorbed onto peptide-conjugated Affi-Gel 10 resin (made using the BioRad protocol), eluted with 0.1 N acetic acid, and neutralized with 2 M Tris-HCl, pH 8. The resulting affinity-purified antibodies were dialyzed into DPBS, concentrated, aliquotted, and stored at -20°C. The antibodies were found to recognize a 125 kDa protein on immunoblots of partially purified A2058 conditioned medium and to preferentially stain some breast carcinoma cells compared to normal breast using immunohistochemical techniques.

An A2058 cDNA library was prepared by purifying poly-A purified mRNA from the cells then size-selecting

- 37 -

° mRNA > 1000 bp for the preparation of cDNA. The cDNA inserts were placed into  $\lambda$ gt11 directionally, using the ProMega cDNA kit using standard methods well-established in the field. LE 392 cells were infected with the  $\lambda$ gt11 and plaques were transferred onto nitrocellulose membranes by overnight incubation at 37°C. The antibody was 5 incubated with the membranes in blocking buffer for 2 hr at room temperature, using approximately twice the concentration of antibody which gave a strong response on Western blot analysis. Secondary antibody was goat anti- 10 rabbit immunoglobulin, and the blot was developed colorimetrically with 4-chloro-1-naphthol.

Positive clones were confirmed by antibody competition with specific peptides but not unrelated peptides. Using this technique and multiple subclonings, 15 we obtained a partial cDNA clone of the autotaxin gene, which we called 4C11. The 4C11 insert was removed from  $\lambda$ gt11 by restriction enzyme digests and subcloned into pBluescript for sequencing by standard Sanger techniques (Sanger, et al., 1977). The 4C11 clone contained bases, 20 including the poly-adenylated tail and the AATAAA adenylation signal locus, i.e., it contained the 3' terminus of the gene. It also included a 627 base open reading frame. Database analysis of this nucleotide sequence revealed that it is unique. The predicated amino 25 acid sequence for 4C11 is 209 amino acids long with exact matches for 7 previously identified ATX peptides: (ATX-20, ATX-34, ATX-102, ATX-104, ATX-204, ATX-215, and ATX-244).

#### EXAMPLE 7

##### Cloning the 5' terminus of ATX

30 Database analysis of the 3' terminus of the ATX gene demonstrated a novel protein. However, we have found an interesting homology that has helped to guide us in exploring its function. ATX had a 45% amino acid identity and a 57% nucleotide identity with PC-1, a marker of B 35 cell activation found on the surface of plasma cells.

- 38 -

Using the PC-1 protein sequence as a guide, we found that ATX peptide homologies were scattered throughout the length of the protein. The only exception was the far amino terminus of PC-1, which includes the transmembrane and intracellular domains, and which had no homologies. Knowing approximate localization of the ATX peptides along the length of ATX, we then amplified different segments of ATX by the PCR (Figure 13). These amplified segments of DNA were then subcloned into plasmids utilizing the TA Cloning kit of ProMega. The PCR amplified DNA could then be sequenced using standard Sanger sequencing techniques (Sanger, et al., 1977).

#### Cloning of full length ATX gene

A reverse transcriptase reaction was performed using total or oligo-(dT) purified RNA from A2058 or N-tera 2D1 cells as template and an anti-sense primer from the 5' end of 4C11 (GCTCAGATAAGGAGGAAAGAG). This was followed by one or two PCR amplification of the resultant cDNA using the commercially available kit from Perkin-Elmer and following manufacturer's directions. These PCR reactions utilized nested antisense primers from 4C11 (GAATCCGTAGGACATCTGCTT and TGTAGGCCAAACAGTTCTGAC) as well as degenerate, nested sense primers deduced from ATX peptides: ATX-101 (AAYTCIATGACARACIGTITTYGTIG and TTYGTIGGITAYGGICCIACITTYAA), ATX-103 (AAYTAYCTIACIAAYGTIGAYGAYAT and GAYGAYATIACICTIGTICCGGIAC), or ATX-224 (TGYTTYGARYTICARGARGCIGGICCCIC). The amplified DNA was then purified from a polyacrylamide gel using standard procedures and ligated into the pCR™ plasmid using the TA cloning kit (Invitrogen Corporation) according to manufacturer's directions.

The 5' RACE kit was utilized to extend the 5' end of ATX cDNA using total RNA from N-tera 2D1 as template and previously obtained sequence as primer (GCTGTCTTCAAACACAGC). The 5' end of the A2058 synthesized

- 39 -

- ° protein was obtained by using previously obtained sequence as primer (CTGGTGGCTGTAATCCATAGC) in a reverse transcriptase reaction with total A2058 RNA as template, followed by PCR amplification utilizing the 5' end of N-tera 2D1 sequence as sense primer
- 5 (CGTGAAGGCAAAGAGAACACG) and a nested antisense primer (GCTGTCTTCAAACACAGC). A2058 DNA encoding ATX is set forth in a SEQ ID NO:68 and the amino acid sequence is provided in SEQ ID NO:69.

10 **DNA sequencing:** DNA sequencing was performed using dideoxy methodology (Sanger, et al. 1977) and (<sup>35</sup>S)dATP (Du Pont, New England Nuclear).

We have found one region between the 5' end of the 4C11 and the ATX peptide designated ATX-101, also referred to as the "hot spot". This region has been

15 sequenced five times with different sequences found each time. The hot spot appears to be located within the region from approximately nucleotide 1670 to 1815. The consensus sequence is represented by amino acids position 559 through 604. Variations found include DNA sequence

20 that results in single and multiple amino acid insertions. One sequence had a stop codon in this region and may have represented an intron. This region has been found to be variable in forms of ATX.

#### EXAMPLE 8

25 Cloning ATX in a human teratocarcinoma cell line

The fact that ATX is present in other cancer cells was confirmed by sequence information from N-tera 2D1, a human teratocarcinoma cell line. For these cells, a prepared cDNA library in λgt10 was amplified and the

30 cDNA inserts were extracted. Using oligonucleotide primers based on known A2058 ATX sequence, DNA segments were amplified by PCR. The DNA segments were then subcloned into plasmids and sequenced as for A2058. We have 3104 bp DNA sequence for N-tera ATX (SEQ ID NO:66)

35 and smaller portions thereof. This includes an open

- 40 -

- ° reading frame that codes for a putative protein containing 861 amino acids (SEQ ID NO:67) and smaller portions thereof. Like the A2058 ATX, the N-tera 2D1 sequence has homologies for multiple ATX peptides (Figure 15). Sequence homology between the A2058 and N-tera 2D1 cells is approximately 99%.

EXAMPLE 9Cloning 5' end of ATX in human normal liver

- The 5' end of ATX has proven difficult to obtain from either tumor cell line to date. Normal human liver mRNA was therefore amplified using the 5' RACE kit (Clontech) with known sequence from A2058 ATX as antisense primer. A DNA segment was obtained and has been sequenced. This segment codes for 979 amino acids, including an initiating methionine (SEQ ID NO:38). The putative protein sequence also includes a 20 amino acid transmembrane domain which is different from the tumor ATX's (SEQ ID NO:54), as shown in Table 7. Both tumorous forms of ATX apparently lack a transmembrane region and are instead secreted proteins.

Table 7

Nucleotide and Amino Acid Sequences Encoding Liver ATX Amino  
Terminus containing the Transmembrane region

## 25 Protein Sequence (SEQ ID NO: 54)

Met Ala Arg Arg Ser Ser Phe Gln Ser Cys Gln Asp Ile Ser Leu Phe Thr Phe Ala Val Gly  
Val Asn Ile Cys Leu Gly Phe Thr Ala His Arg Ile Lys Arg Ala Glu Gly Trp

## 30 DNA Sequence (SEQ ID NO: 53)

ATGGCAAGGA GGAGCTCGTT CCAGTCGTGT CAAGATATAT CCCTGTTTAC  
TTTTGCCGTT GGAGTCAATA TCTGCTTAGG ATTCAGTGCA CATCGAATTA  
AGAGAGCAGA AGGATGG

35

- 41 -

EXAMPLE 10Domains of ATX

Searches of protein databases (Pearson, et. al. 1988) confirmed that the homology between ATX and PC-1 was present throughout the length of the extracellular portion of the molecules (Buckley, et. al., 1990; Funakoshi, et. al. 1992). There is a 45% amino acid identity and a 64% similarity between the 2 protein sequences (Fig. 18). For the cDNA sequence, the identity is ~57%.

These proteins share several interesting properties and domains (Fig. 19). Both have a number of potential N-linked glycosylation sites: four for ATX (Asn54, Asn463, Asn577, Asn859) and nine for PC-1. Both have adjacent somatomedin B domains near the amino end of the extracellular domain. This somatomedin B domain is a cysteine-rich region containing 3 presumed cystine cross-linkages. ATX has 33 Cys residues and PC-1 has 37; 30 of these Cys residues are identical in placement. Both proteins also contain the loop region of an EF hand (Buckley, et. al. 1990; Kretsinger, 1987). In addition, both proteins have a transmembrane/signal peptide region with a short intracellular peptide, common in ectoenzymes (Maroux, 1987). However, the amino acid identity between ATX and PC-1 in the intracellular and transmembrane regions is only 11%.

Finally, both proteins have a region homologous to the bovine intestinal phosphodiesterase enzymatic domain with conservation of the threonine that is thought to act as the intermediate phosphate binding site (Culp, et al. 1985). PC-1 has been demonstrated to have phosphodiesterase type I, nucleotide pyrophosphatase, and threonine-specific kinase enzymatic activities (Rebbe, et al. 1991; Oda, et al. 1991). In order to test whether purified ATX had type I phosphodiesterase activity, samples were incubated with p-nitrophenyl thymidine-5'-monophosphate at pH 8.9 for 30 min. Samples were assayed

- 42 -

- ° in a 100  $\mu$ l volume containing 50 mM Tris-HCl, pH 8.9 and 5 mM p-nitrophenyl thymidine-5'-monophosphate. After a 30 minute incubation at 37 °C the reactions were terminated by addition of 900  $\mu$ l 0.1 N NaOH and the amount of product formed was determined by reading the absorbance at 410 nm.
- 5 ATX was found to hydrolyze the p-nitrophenyl thymidine-5'-monophosphate (Razzell, 1963) at a rate of 10 pmol/ng/min, a reaction rate similar to that reported for PC-1 (Oda, et al. 1993).

#### REFERENCES

- 10 Atnip, K.D., et al. (1987) *Biochem. Biophys. Res. Comm.* 146, 996-1002
- Buckley, M.F., Loveland, K.A., McKinsty, W.J., Garson, O. M. and Goding, J.W. (1990) *J. Biol. Chem.* 265, 17506-17511
- 15 Culp, J.S., Blytt, H.J., Hermodson, M. and Butler, L.G. (1985) *J. Biol. Chem.* 260, 8320-8324
- Dennis, J. W., Koch, K., Yousefi, S. and VanderElst, I. 20 (1990) *Cancer Res.* 50, 1867-1872
- Funakoshi, I., Kato, H., Horie, K., Yano, T., Hori, Y., Kobayashi, H., Inoue, T., Suzuki, H., Fukui, S., Tsukahara, M., Kajii, T. and Yamashina, I. (1992) *Arch. Biochem. Biophys.* 295, 180-187
- 25 Gospodarowicz, D. (1984) *Proc. Natl. Acad. Sci. USA* 81, 6963-6967
- 30 Guirguis, R., et al. (1987) *Nature* 329, 261-263
- Jouanneau, J., et al. (1991) *Proc. Natl. Acad. Sci. USA* 88, 2893-2897
- 35 Kahan, B.W. et al., (1987) *Cancer Res.* 47, 6324-6328



- 43 -

- ° Kretsinger, R.H. (1987) *Cold Spring Harbor Symp. Quant. Bio.* 52, 499-510
- Kohler and Milstein, (1975) *Nature* 256:495-497
- 5 Kohn, E.C., et al. (1990) *Int. J. Cancer* 46, 287-292  
Laemmli U.K. (1970) *Nature* 227, 680-685
- Landsteiner, *Specificity of Serological Reactions* (Dover Publications, New York, 1962)
- 10
- Liotta, L.A., et al. (1986) *Proc. Natl. Acad. Sci. USA* 83, 3302-3306
- 15 Liotta, L.A., et al. (1988) *Cancer Surveys* 7, 631-652
- Maciag, T., et al. (1984) *Sci.* 225, 932-935
- Maroux, S. (1987) In A.J. Kenny and A.J. Turner (eds.)  
20 *Mammalian Ectoenzymes*, Elsevier Science Publishers B.V., Amsterdam, The Netherlands, 15-45
- McCarthy, J.B., et al. (1984) *J. Cell Biol.* 98, 1474-1480  
Microbiology, Hoeber Medical Division (Harper and Row,  
25 1969)
- Nabi, I.R., et al. (1990) *Cancer Res.* 50, 409-414
- Neuhoff, V., et al. (1988) *Electrophoresis* 9, 255-262
- 30 Oda, Y., Kuo, M.-D., Huang, S.S. and Huang, J.S. (1991) *J. Biol. Chem.* 266, 16791-16795
- Oda, Y., Kuo, M.-D., Huang, S.S. and Huang, J.S. (1993) *J. Biol. Chem.* 268, 27318-27326  
35

- 44 -

- ° O'Farrell, P.H. (1975) *J. Biol. Chem.* 250, 4007-4021.
- O'Farrell, P.H., et al. (1977) *Cell* 12, 1133-1142
- 5 Ogier-Denis, E., Trugnan, G., Sapin, C., Aubery, M. and  
Codogno, P. (1990) *J. Biol. Chem.* 265, 5366-5369
- Ohnishi, T., et al. (1990) *J. Neurosurg.* 73, 881-888
- 10 Pearson, W.R., et al. (1988) *Proc. Natl. Acad. Sci. USA*  
85, 2444-2448
- Razzell, W.E. (1963) *Methods Enzymol.* 6, 236-258
- 15 Rebbe, N., F., Tong, B.D., Finley, E.M. and Hickman, S.  
(1991) *Proc. Natl. Acad. Sci. USA* 88, 5192-5196
- Rosen, E.M. et al., (1989) *In Vitro Cell Devel. Biol.* 25,  
163-173
- 20 Rosen, E.M., et al. (1990) *Proc. Soc. Exp. Biol. Med.* 195,  
34-43
- Ruff, M., et al. (1985) *Clin. Immunol. Immunopath.* 37,  
387-396
- 25 Sanger, F. et al. (1977) *Proc. National Acad. Sci. USA.*  
74, 5463-5467
- Schnor, S.L., et al. (1988) *J. Cell Sci.* 90, 391-399
- 30 Seftor, R. E. B., Seftor, E. A., Grimes, W. J., Liotta, L.  
A.,  
Stetler-Stevenson, W. G., Welch, D. R. and Hendrix, M. J.  
C. (1991) *Melanoma Res.* 1, 43-54
- 35 Silletti, S., et al. (1991) *Cancer Res.* 51, 3507-3511

- 45 -

- ° Singer, S.J. and Kupfer, A. (1986) *Ann. Rev. Cell Biol.* 2, 337-365
- Stites et al., editors, Basic and Clinical Immunology,  
(Lange Medical Publications, Los Altos, CA, Fourth  
5 edition)
- Stoker, M., et al. (1987) *Nature* 327, 239-242
- Stone, M, et al. (1989) A Practical Guide to Protein and  
10 Peptide Purification for Microsequencing (Matsudaira,  
P.T., ed.) pgs. 33-47, Academic Press, N.Y.
- Stracke, M.L. et al., *Biochem. Biophys. Res. Comm.* 153,  
1076-1083
- 15 Stracke, M.L., et al. (1978) *Biochem. Biophys. Res. Comm.*  
146, 339-345
- Stracke, M.L., et al. (1987) *Biochem. Biophys. Res. Comm.*  
20 147, 339-345
- Stracke, M.L., et al. (1988) *Biochem. Biophys. Res. Comm.*  
153, 1076-1083
- 25 Tamm, I., et al., (1989) *J. Exp. Med.* 170, 1649-1669
- Taraboletti, G., (1987) *J. Cell Biol.* 105, 2409-2415
- Todaro, G.J., et al. (1980) *Proc. Natl. Acad. Sci. USA* 77,  
30 5258-5262
- Van Snick, J. (1990) *Ann. Rev. Immunol.* 8, 253-278
- Wang, J.M., et al. (1990) *Biochem. Biophys. Res. Comm.*  
35 169, 165-170

- 46 -

- ° Watanabe, H., et al. (1990) *J. Cell Biol.* **111**, 2097-2108
- Watanabe, H., et al. (1991) *J. Biol. Chem.* **266**, 13442-13448
- 5 Weidner, K.M., et al. (1990) *J. Cell. Biol.* **111**, 2097-2108
- Williams et al., Methods in Immunology and Immunochimistry, Vol. 1 (Academic Press, New York, 1967)
- 10 Yoshimura, T. (1987) *Proc. Natl. Acad. Sci. USA* **84**, 9233-9237

\* \* \* \* \*

All publications mentioned hereinabove are hereby incorporated in their entirety by reference.

- 15 While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be appreciated by one skilled in the art from a reading of this disclosure that various changes in form and detail can be made without departing from the true
- 20 scope of the present invention and appended claims.

25

30

35

- 47 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: STRACKE, MARY; LIOTTA, LANCE;  
SCHIFFMANN, ELLIOTT; KRUTZSCH,  
HENRY; MURATA, JUN
- 5 (ii) TITLE OF INVENTION: MOTILITY STIMULATING  
PROTEIN USEFUL IN CANCER DIAGNOSIS AND  
THERAPY
- (iii) NUMBER OF SEQUENCES: 69
- 10 (iv) CORRESPONDENCE ADDRESS:  
(A) ADDRESSEE: MORGAN & FINNEGAN  
(B) STREET: 345 PARK AVENUE  
(C) CITY: NEW YORK  
(D) STATE: NEW YORK  
(E) COUNTRY: U.S.A.  
(F) ZIP: 10154
- 15 (v) COMPUTER READABLE FORM:  
(A) MEDIUM TYPE: Floppy Disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: WordPerfect 5.1
- (vi) CURRENT APPLICATION DATA:  
20 (A) APPLICATION NUMBER:  
(B) FILING DATE: 24-MAY-1995  
(C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA  
(A) APPLICATION NUMBER: 08/346,455  
(B) FILING DATE: 28-NOV-1994
- 25 (vii) PRIOR APPLICATION DATA  
(A) APPLICATION NUMBER: 08/249,182  
(B) FILING DATE: 25-MAY-1994
- (vii) PRIOR APPLICATION DATA  
(A) APPLICATION NUMBER: 07/822,043  
(B) FILING DATE: 17-JAN-1992
- 30 (viii) ATTORNEY/AGENT INFORMATION:  
(A) NAME: DOROTHY R. AUTH  
(B) REGISTRATION NUMBER: 36,434  
(C) DOCKET NUMBER: 2026-4149US2
- (ix) TELECOMMUNICATION INFORMATION:  
35 (A) TELEPHONE: (212) 758-4800  
(B) TELEFAX: (212) 751-6849

- 48 -

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Trp His Val Ala Arg  
1 5

(2) INFORMATION FOR SEQ ID NO:2:

10

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

15

Pro Leu Asp Val Tyr Lys  
1 5

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

20

- (A) LENGTH: 5
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Tyr Pro Ala Phe Lys  
1 5

25

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Gln Ala Glu Val Ser  
1 5

(2) INFORMATION FOR SEQ ID NO:5:

35

- 49 -

- ° (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 10  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:
- 5 Pro Glu Glu Val Thr Arg Pro Asn Tyr Leu  
1 5 10
- (2) INFORMATION FOR SEQ ID NO:6:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 9  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear
- 10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:
- Tyr Asp Val Pro Trp Asn Glu Thr Ile  
1 5
- 15 (2) INFORMATION FOR SEQ ID NO:7:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 10  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear
- 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:
- Val Pro Pro Phe Glu Asn Ile Glu Leu Tyr  
1 5 10
- (2) INFORMATION FOR SEQ ID NO:8:
- 25 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 11  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:
- 30 Gly Gly Gln Pro Leu Trp Ile Thr Ala Thr Lys  
1 5 10
- (2) INFORMATION FOR SEQ ID NO:9:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 16
- 35

- 50 -

- (B) TYPE: amino acid  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Val Asn Ser Met Gln Thr Val Phe Val Gly Tyr Gly  
1 5 10

5

Pro Thr Phe Lys  
15

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 12  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Asp Ile Glu His Leu Thr Ser Leu Asp Phe Phe Arg  
1 5 10

15

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

20

- (A) LENGTH: 23  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Thr Glu Phe Leu Ser Asn Tyr Leu Thr Asn Val Asp  
1 5 10

25

Asp Ile Thr Leu Val Pro Glu Thr Leu Gly Arg  
15 20

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

30

- (A) LENGTH: 18  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

35

GTTGGCAGCN ACRTGCCA

18



- 51 -

## (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 18  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TGGCAYGTNG CTGCCAAC

18

## (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CTTGAAGGCA GGGTA

15

## (2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

TAYCCTGCNT TYAAG

15

## (2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GGTNACYTCY TCAGG

15

- 52 -

- 5 (2) INFORMATION FOR SEQ ID NO:17:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 15
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:
- CCTGARGARG TNACC 15
- 10 (2) INFORMATION FOR SEQ ID NO:18:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 21
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- 15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:
- NGTNGCRTCR AATGGCACRT C 21
- 20 (2) INFORMATION FOR SEQ ID NO:19:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 21
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:
- GAYGTGCCAT TYGAYGCNAC N 21
- (2) INFORMATION FOR SEQ ID NO:20:
- 30 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 21
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:
- 35 GTTDA TRTTS TCRAATGGGG G 21

- 53 -

## (2) INFORMATION FOR SEQ ID NO:21:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

CCCCCATTG AGAACATCAA C

21

## 10 (2) INFORMATION FOR SEQ ID NO:22:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## 15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

CTTNGTNGCN GTDATCCANA RGGGYTGGCC GCC

33

## 20 (2) INFORMATION FOR SEQ ID NO:23:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GGCGGCCARC CCYTNTGGAT HACNGCNACN AAG

33

## (2) INFORMATION FOR SEQ ID NO:24:

## 30 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

35

- 54 -

° (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

CTTRAAGGTG GGGCCRTAGC CCACRAAGAC TGTYTG CAT

39

(2) INFORMATION FOR SEQ ID NO:25:

- 5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 39  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

ATGCARACAG TCTTYGTGGG CTAYGGCCCC ACCTTYAAR

39

(2) INFORMATION FOR SEQ ID NO:26:

- 15 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 9  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

20 Gln Tyr Leu His Gln Tyr Gly Ser Ser  
1 5

(2) INFORMATION FOR SEQ ID NO:27:

- 25 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 5  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

30 Val Leu Asn Tyr Phe  
1 5

(2) INFORMATION FOR SEQ ID NO:28:

- 35 (i) SEQUENCE CHARACTERISTICS:

- 55 -

- (A) LENGTH: 5
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Tyr Leu Asn Ala Thr  
1 5

(2) INFORMATION FOR SEQ ID NO:29:

- 10 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 11
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

His Leu Leu Tyr Gly Arg Pro Ala Val Leu Tyr  
1 5 10

(2) INFORMATION FOR SEQ ID NO:30:

- 20 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 11
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Ser Tyr Pro Glu Ile Leu Thr Pro Ala Asp Asn  
1 5 10

(2) INFORMATION FOR SEQ ID NO:31:

- 30 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 14
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

35

- 56 -

° (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Xaa Tyr Gly Phe Leu Phe Pro Pro Tyr Leu Ser Ser  
 1 5 10  
 Ser Pro

5 (2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 13  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Thr Phe Pro Asn Leu Tyr Thr Phe Ala Thr Gly Leu  
 1 5 10  
 Tyr

15

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 22  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Val Asn Val Ile Ser Gly Pro Ile Asp Asp Tyr Asp  
 1 5 10  
 Tyr Asp Gly Leu His Asp Thr Glu Asp Lys  
 15 20

25

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 829  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: Unknown

30

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: No

(vi) ORIGINAL SOURCE:

35

- 57 -

- 5 (A) ORGANISM: Human  
 (B) STRAIN:  
 (C) INDIVIDUAL ISOLATE:  
 (D) DEVELOPMENTAL STAGE:  
 (E) HAPLOTYPE:  
 (F) TISSUE TYPE:  
 (G) CELL TYPE: Melanoma  
 (H) CELL LINE: A2058  
 (I) ORGANELLE:

- 10 (ix) FEATURE:  
 (A) NAME/KEY:  
 (B) LOCATION:  
 (C) IDENTIFICATION METHOD:  
 (D) OTHER INFORMATION: Putative protein  
 sequence of A2058 Autotaxin

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

|    |     |     |     |     |     |     |     |     |     |     |     |     |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
|    | Cys | His | Asp | Phe | Asp | Glu | Leu | Cys | Leu | Lys | Thr | Ala |
|    | 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |
| 15 | Arg | Gly | Trp | Glu | Cys | Thr | Lys | Asp | Arg | Cys | Gly | Glu |
|    |     | 15  |     |     |     |     |     | 20  |     |     |     |     |
|    | Val | Arg | Asn | Glu | Glu | Asn | Ala | Cys | His | Cys | Ser | Glu |
|    | 25  |     |     |     |     | 30  |     |     |     |     | 35  |     |
|    | Asp | Cys | Leu | Ala | Arg | Gly | Asp | Cys | Cys | Thr | Asn | Tyr |
|    |     |     | 40  |     |     |     |     |     | 45  |     |     |     |
|    | Gln | Val | Val | Cys | Lys | Gly | Glu | Ser | His | Trp | Val | Asp |
|    | 50  |     |     |     |     | 55  |     |     |     |     | 60  |     |
| 20 | Asp | Asp | Cys | Glu | Glu | Ile | Lys | Ala | Ala | Glu | Cys | Pro |
|    |     |     |     | 65  |     |     |     |     |     | 70  |     |     |
|    | Ala | Gly | Phe | Val | Arg | Pro | Pro | Leu | Ile | Ile | Phe | Ser |
|    |     | 75  |     |     |     |     | 80  |     |     |     |     |     |
|    | Val | Asp | Gly | Phe | Arg | Ala | Ser | Tyr | Met | Lys | Lys | Gly |
|    | 85  |     |     |     | 90  |     |     |     |     | 95  |     |     |
|    | Ser | Lys | Val | Met | Pro | Asn | Ile | Glu | Lys | Leu | Arg | Ser |
|    |     |     | 100 |     |     |     |     |     | 105 |     |     |     |
| 25 | Cys | Gly | Thr | His | Ser | Pro | Tyr | Met | Arg | Pro | Val | Tyr |
|    | 110 |     |     |     |     | 115 |     |     |     |     | 120 |     |
|    | Pro | Thr | Lys | Thr | Phe | Pro | Asn | Leu | Tyr | Thr | Leu | Ala |
|    |     |     |     | 125 |     |     |     |     |     | 130 |     |     |
|    | Thr | Gly | Leu | Tyr | Pro | Glu | Ser | His | Gly | Ile | Val | Gly |
|    |     | 135 |     |     |     | 140 |     |     |     |     |     |     |
|    | Asn | Ser | Met | Tyr | Asp | Pro | Val | Phe | Asp | Ala | Thr | Phe |
| 30 | 145 |     |     |     | 150 |     |     |     |     | 155 |     |     |
|    | His | Leu | Arg | Gly | Arg | Glu | Lys | Phe | Asn | His | Arg | Trp |
|    |     |     | 160 |     |     |     |     | 165 |     |     |     |     |
|    | Trp | Gly | Gly | Gln | Pro | Leu | Trp | Ile | Thr | Ala | Thr | Lys |
|    | 170 |     |     |     |     | 175 |     |     |     |     | 180 |     |
|    | Gln | Gly | Val | Lys | Ala | Gly | Thr | Phe | Phe | Trp | Ser | Val |
| 35 |     |     |     | 185 |     |     |     |     |     | 190 |     |     |

- 58 -

° Val Ile Pro His Glu Arg Arg Ile Leu Thr Ile Leu  
195 200  
Arg Trp Leu Thr Leu Pro Asp His Glu Arg Pro Ser  
205 210 215  
Val Tyr Ala Phe Tyr Ser Glu Gln Pro Asp Phe Ser  
220 225  
5 Gly His Lys Tyr Gly Pro Phe Gly Pro Glu Glu Ser  
230 235 240  
Ser Tyr Gly Ser Pro Phe Thr Pro Ala Lys Arg Pro  
245 250  
Lys Arg Lys Val Ala Pro Lys Arg Arg Gln Glu Arg  
255 260  
Pro Val Ala Pro Pro Lys Lys Arg Arg Arg Lys Ile  
265 270 275  
10 His Arg Met Asp His Tyr Ala Ala Glu Thr Arg Gln  
280 285  
Asp Lys Met Thr Asn Pro Leu Arg Glu Ile Asp Lys  
290 295 300  
Ile Val Gly Gln Leu Met Asp Gly Leu Lys Gln Leu  
305 310  
Lys Leu Arg Arg Cys Val Asn Val Ile Phe Val Gly  
315 320  
15 Asp His Gly Met Glu Asp Val Thr Cys Asp Arg Thr  
325 330 335  
Glu Phe Leu Ser Asn Tyr Leu Thr Asn Val Asp Asp  
340 345  
Ile Thr Leu Val Pro Gly Thr Leu Gly Arg Ile Arg  
350 355 360  
Ser Lys Phe Ser Asn Asn Ala Lys Tyr Asp Pro Lys  
365 370  
20 Ala Ile Ile Ala Asn Leu Thr Cys Lys Lys Pro Asp  
375 380  
Gln His Phe Lys Pro Tyr Leu Lys Gln His Leu Pro  
385 390 395  
Lys Arg Leu His Tyr Ala Asn Asn Arg Arg Ile Glu  
400 405  
25 Asp Ile His Leu Leu Val Glu Arg Arg Trp His Val  
410 415 420  
Ala Arg Lys Pro Leu Asp Val Tyr Lys Lys Pro Ser  
425 430  
Gly Lys Cys Phe Phe Gln Gly Asp His Gly Phe Asp  
435 440  
Asn Lys Val Asn Ser Met Gln Thr Val Phe Val Gly  
445 450 455  
30 Tyr Gly Pro Thr Phe Lys Tyr Lys Thr Lys Val Pro  
460 465  
Pro Phe Glu Asn Ile Glu Leu Tyr Asn Val Met Cys  
470 475 480  
Asp Leu Leu Gly Leu Lys Pro Ala Pro Asn Asn Gly  
485 490  
Thr His Gly Ser Leu Asn His Leu Leu Arg Thr Asn  
495 500  
35 Thr Phe Arg Pro Thr Met Pro Glu Glu Val Thr Arg  
505 510 515



- 59 -

|    |     |     |     |     |     |     |     |     |     |     |     |     |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
|    | Pro | Asn | Tyr | 520 | Gly | Ile | Met | Tyr | Leu | Gln | Ser | Asp |
|    | Asp | Asp | Leu | 530 | Gly | Cys | Thr | Cys | Asp | Asp | Lys | Val |
|    | Pro | Lys | Asn | 545 | Lys | Leu | Asp | Glu | Leu | Asn | Lys | Arg |
| 5  | His | Thr | Lys | 555 | Gly | Ser | Thr | Glu | Glu | Arg | His | Leu |
|    | Tyr | Gly | Arg | 565 | Pro | Ala | Val | Leu | Tyr | Arg | Thr | Arg |
|    | Asp | Ile | Leu | 580 | Tyr | His | Thr | Asp | Phe | Glu | Ser | Gly |
|    | Ser | Glu | Ile | 590 | Phe | Leu | Met | Leu | Leu | Trp | Thr | Ser |
| 10 | Thr | Val | Ser | 605 | Lys | Gln | Ala | Glu | Val | Ser | Ser | Val |
|    | Asp | His | Leu | 615 | Thr | Ser | Cys | Val | Arg | Pro | Asp | Val |
|    | Val | Ser | Pro | 625 | Ser | Phe | Ser | Gln | Asn | Cys | Leu | Ala |
|    | Lys | Asn | Asp | 640 | Lys | Gln | Met | Ser | Tyr | Gly | Phe | Leu |
| 15 | Pro | Pro | Tyr | 650 | Leu | Ser | Ser | Ser | Pro | Glu | Ala | Lys |
|    | Asp | Ala | Phe | 665 | Leu | Val | Thr | Asn | Met | Val | Pro | Met |
|    | Pro | Ala | Phe | 675 | Lys | Arg | Val | Trp | Asn | Tyr | Phe | Gln |
|    | Val | Leu | Val | 685 | Lys | Lys | Tyr | Ala | Ser | Glu | Arg | Asn |
| 20 | Val | Asn | Val | 700 | Ile | Ser | Gly | Pro | Ile | Phe | Asp | Tyr |
|    | Tyr | Asp | Gly | 710 | Leu | His | Asp | Thr | Glu | Asp | Lys | Ile |
|    | Gln | Tyr | Val | 725 | Glu | Gly | Ser | Ser | Ile | Pro | Val | Pro |
| 25 | His | Tyr | Tyr | 735 | Ser | Ile | Ile | Thr | Ser | Cys | Leu | Asp |
|    | Thr | Gln | Pro | 745 | Ala | Asp | Lys | Cys | Asp | Gly | Pro | Leu |
|    | Val | Ser | Ser | 760 | Phe | Ile | Leu | Pro | His | Arg | Pro | Asp |
|    | Glu | Glu | Ser | 770 | Cys | Asn | Ser | Ser | Glu | Asp | Glu | Ser |
| 30 | Trp | Val | Glu | 785 | Glu | Leu | Met | Lys | Met | His | Thr | Ala |
|    | Val | Arg | Asp | 795 | Ile | Glu | His | Leu | Thr | Ser | Leu | Asp |
|    | Phe | Arg | Lys | 805 | Thr | Ser | Arg | Ser | Tyr | Pro | Glu | Ile |
| 35 | Thr | Leu | Lys | 820 | Thr | Tyr | Leu | His | Thr | Tyr | Glu | Ser |

- 60 -

Ile

## (2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
- 5 (A) LENGTH: 2946
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: Unknown
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: No
- 10 (vi) ORIGINAL SOURCE:
- (A) ORGANISM: Human
- (B) STRAIN:
- (C) INDIVIDUAL ISOLATE:
- (D) DEVELOPMENTAL STAGE:
- (E) HAPLOTYPE:
- 15 (F) TISSUE TYPE:
- (G) CELL TYPE: Melanoma
- (H) CELL LINE: A2058
- (I) ORGANELLE:
- (ix) FEATURE:
- (A) NAME/KEY:
- (B) LOCATION:
- 20 (C) IDENTIFICATION METHOD:
- (D) OTHER INFORMATION: Partial DNA Sequence of A2058 Autotaxin
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:
- |    |             |            |            |            |     |
|----|-------------|------------|------------|------------|-----|
|    | GCTGCCATGA  | CTTTGATGAG | CTGTGTTTGA | AGACAGCCCG | 40  |
|    | TGGCTGGGAG  | TGTACTAAGG | ACAGATGTGG | AGAAGTCAGA | 80  |
| 25 | AATGAAGAAA  | ATGCCTGTCA | CTGCTCAGAG | GACTGCTTGG | 120 |
|    | CCAGGGGAGA  | CTGCTGTACC | AATTACCAAG | TGGTTTGCAA | 160 |
|    | AGGAGAGTCG  | CATTGGGTTG | ATGATGACTG | TGAGGAAATA | 200 |
|    | AAGGCCGCGAG | AATGCCCTGC | AGGGTTTGTG | CGCCCTCCAT | 240 |
|    | TAATCATCTT  | CTCCGTGGAT | GGCTTCCGTG | CATCATACAT | 280 |
|    | GAAGAAAGGC  | AGCAAAGTCA | TGCCTAATAT | TGAAAACTA  | 320 |
|    | AGGTCTTGTG  | GCACACACTC | TCCCTACATG | AGGCCGGTGT | 360 |
| 30 | ACCCAACTAA  | AACCTTTCCT | AACTTATACA | CTTTGGCCAC | 400 |
|    | TGGGCTATAT  | CCAGAATCAC | ATGGAATTGT | TGGCAATTCA | 440 |
|    | ATGTATGATC  | CTGTATTTGA | TGCCACTTTT | CATCTGCGAG | 480 |
|    | GGCGAGAGAA  | ATTTAATCAT | AGATGGTGGG | GAGGTCAACC | 520 |
|    | GCTATGGATT  | ACAGCCACCA | AGCAAGGGGT | GAAAGCTGGA | 560 |
|    | ACATTCTTTT  | GGTCTGTTGT | CATCCCTCAC | GAGCGGAGAA | 600 |
|    | TATTAACCAT  | ATTGCGGTGG | CTCACCCTGC | CAGATCATGA | 640 |
|    | GAGGCCTTCG  | GTCTATGCCT | TCTATTCTGA | GCAACCTGAT | 680 |
| 35 | TTCTCTGGAC  | ACAAATATGG | CCCTTTCGGC | CCTGAGGAGA | 720 |

- 61 -

|    |            |            |             |             |      |
|----|------------|------------|-------------|-------------|------|
|    | GTAGTTATGG | CTCACCTTTT | ACTCCGGCTA  | AGAGACCTAA  | 760  |
|    | GAGGAAAGTT | GCCCCTAAGA | GGAGACAGGA  | AAGACCAGTT  | 800  |
|    | GCTCCTCCAA | AGAAAAGAAG | AAGAAAAATA  | CATAGGATGG  | 840  |
|    | ATCATTATGC | TGCGGAAACT | CGTCAGGACA  | AAATGACAAA  | 880  |
|    | TCCTCTGAGG | GAAATCGACA | AAATTGTGGG  | GCAATTAATG  | 920  |
|    | GATGGACTGA | AACAACATAA | ACTGCGTCGG  | TGTGTCAACG  | 960  |
| 5  | TCATCTTTGT | CGGAGACCAT | GGAATGGAAG  | ATGTCACATG  | 1000 |
|    | TGATAGAACT | GAGTTCTTGA | GTAATTACCT  | AACTAATGTG  | 1040 |
|    | GATGATATTA | CTTTAGTGCC | TGGAACCTCTA | GGAAGAATTC  | 1080 |
|    | GATCCAAATT | TAGCAACAAT | GCTAAATATG  | ACCCCAAAGC  | 1120 |
|    | CATTATTGCC | AATCTCACGT | GTAAAAAACC  | AGATCAGCAC  | 1160 |
|    | TTTAAGCCTT | ACTTGAAACA | GCACCTTCCC  | AAACGTTTGC  | 1200 |
|    | ACTATGCCAA | CAACAGAAGA | ATTGAGGATA  | TCCATTTATT  | 1240 |
|    | GGTGGAACGC | AGATGGCATG | TTGCAAGGAA  | ACCTTTGGAT  | 1280 |
| 10 | GTTTATAAGA | AACCATCAGG | AAAATGCTTT  | TTCCAGGGAG  | 1320 |
|    | ACCACGGATT | TGATAACAAG | GTCAACAGCA  | TGCAGACTGT  | 1360 |
|    | TTTTGTAGGT | TATGGCCCAA | CATTTAAGTA  | CAAGACTAAA  | 1400 |
|    | GTGCCTCCAT | TTGAAAACAT | TGAACCTTAC  | AATGTTATGT  | 1440 |
|    | GTGATCTCCT | GGGATTGAAG | CCAGCTCCTA  | ATAATGGGAC  | 1480 |
|    | CCATGGAAGT | TTGAATCATC | TCCTGCGCAC  | TAATACCTTC  | 1520 |
|    | AGGCCAACCA | TGCCAGAGGA | AGTTACCAGA  | CCCAATTATC  | 1560 |
|    | CAGGGATTAT | GTACCTTCAG | TCTGATTTTG  | ACCTGGGCTG  | 1600 |
| 15 | CACTTGTGAT | GATAAGGTAG | AGCCAAAGAA  | CAAGTTGGAT  | 1640 |
|    | GAAGTCAACA | AACGGCTTCA | TACAAAAGGG  | TCTACAGAAG  | 1680 |
|    | AGAGACACCT | CCTCTATGGG | CGACCTGCAG  | TGCTTTATCG  | 1720 |
|    | GACTAGATAT | GATATCTTAT | ATCACACTGA  | CTTTGAAAGT  | 1760 |
|    | GGTTATAGTG | AAATATTCCT | AATGCTACTC  | TGGACATCAT  | 1800 |
|    | ATACTGTTTC | CAAACAGGCT | GAGGTTTCCA  | GCGTTCCTGA  | 1840 |
|    | CCATCTGACC | AGTTGCGTCC | GGCCTGATGT  | CCGTGTTTCT  | 1880 |
| 20 | CCGAGTTTCA | GTCAGAACTG | TTTGGCCTAC  | AAAAATGATA  | 1920 |
|    | AGCAGATGTC | CTACGGATTG | CTCTTTCCCTC | CTTATCTGAG  | 1960 |
|    | CTCTTCACCA | GAGGCTAAAT | ATGATGCATT  | CCTTGTAACC  | 2000 |
|    | AATATGGTTC | CAATGTATCC | TGCTTTCAAA  | CGGGTCTGGA  | 2040 |
|    | ATTATTTCCA | AAGGGTATTG | GTGAAGAAAT  | ATGCTTCGGA  | 2080 |
|    | AAGAAATGGA | GTAAACGTGA | TAAGTGGACC  | AATCTTCGAC  | 2120 |
|    | TATGACTATG | ATGGCTTACA | TGACACAGAA  | GACAAAATAA  | 2160 |
|    | ACAGTACGT  | GGAAGGCAGT | TCCATTCCCTG | TTCCAACTCA  | 2200 |
| 25 | CTACTACAGC | ATCATCACCA | GCTGTCTGGA  | TTTCACTCAG  | 2240 |
|    | CCTGCCGACA | AGTGTGACGG | CCCTCTCTCT  | GTGTCCTCCT  | 2280 |
|    | TCATCCTGCC | TCACCGGCCT | GACAAAGAGG  | AGAGCTGCAA  | 2320 |
|    | TAGCTCAGAG | GACGAATCAA | AATGGGTAGA  | AGAAGCTCATG | 2360 |
|    | AAGATGCACA | CAGCTAGGGT | GCGTGACATT  | GAACATCTCA  | 2400 |
|    | CCAGCCTGGA | CTTCTTCCGA | AAGACCAGCC  | GCAGCTACCC  | 2440 |
|    | AGAAATCCTG | ACACTCAAGA | CATACCTGCA  | TACATATGAG  | 2480 |
| 30 | AGCGAGATTT | AACTTTCTGA | GCATCTGCAG  | TACAGTCTTA  | 2520 |
|    | TCAACTGGTT | GTATATTTTT | ATATTGTTTT  | TGTATTTATT  | 2560 |
|    | AATTTGAAAC | CAGGACATTA | AAAATGTTAG  | TATTTTAATC  | 2600 |
|    | CTGTACCAA  | TCTGACATAT | TATGCCTGAA  | TGACTCCACT  | 2640 |
|    | GTTTTTCTCT | AATGCTTGAT | TTAGGTAGCC  | TTGTGTTCTG  | 2680 |
|    | AGTAGAGCTT | GTAATAAATA | CTGCAGCTTG  | AGAAAAAGTG  | 2720 |
|    | GAAGCTTCTA | AATGGTGCTG | CAGATTTGAT  | ATTTGCATTG  | 2760 |
|    | AGGAAATATT | AATTTTCCAA | TGCACAGTTG  | CCACATTTAG  | 2800 |
| 35 | TCCTGTACTG | TATGGAAACA | CTGATTTTGT  | AAAGTTGCCT  | 2840 |

- 62 -

TTATTTGCTG TTAAGTGTGA ACTATGACAG ATATATTTAA 2880  
 GCCTTATAAA CCAATCTTAA ACATAATAAA TCACACATTC 2920  
 AGTTTATAAA AAAAAAAAAA AAAAAA 2946

## (2) INFORMATION FOR SEQ ID NO:36:

- 5 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 788  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: Unknown
- (ii) MOLECULE TYPE: protein
- 10 (iii) HYPOTHETICAL: No
- (vi) ORIGINAL SOURCE:  
 (A) ORGANISM: Human  
 (B) STRAIN:  
 (C) INDIVIDUAL ISOLATE:  
 (D) DEVELOPMENTAL STAGE:  
 15 (E) HAPLOTYPE:  
 (F) TISSUE TYPE:  
 (G) CELL TYPE: teratocarcinoma  
 (H) CELL LINE: N-tera 2D1  
 (I) ORGANELLE:
- (ix) FEATURE:  
 20 (A) NAME/KEY:  
 (B) LOCATION:  
 (C) IDENTIFICATION METHOD:  
 (D) OTHER INFORMATION: N-tera 2D1 putative  
 ATX protein sequence
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:
- 25 Cys Asp Asn Leu Cys Lys Ser Tyr Thr Ser Cys Cys  
 1 5 10  
 His Asp Phe Asp Glu Leu Cys Leu Lys Thr Ala Arg  
 15 20  
 Ala Trp Glu Cys Thr Lys Asp Arg Cys Gly Glu Val  
 25 30 35  
 Arg Asn Glu Glu Asn Ala Cys His Cys Ser Glu Asp  
 40 45  
 30 Cys Leu Ala Arg Gly Asp Cys Cys Thr Asn Tyr Gln  
 50 55 60  
 Val Val Cys Lys Gly Glu Ser His Trp Val Asp Asp  
 65 70  
 Asp Cys Glu Glu Ile Lys Ala Ala Glu Cys Leu Gln  
 75 80  
 Val Asp Ser Pro Ser Ile Asn His Leu Leu Arg Gly  
 85 90 95
- 35

|    |     |     |     |     |     |     |     |     |     |     |     |     |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 0  | Trp | Leu | Pro | Met | Thr | Ser | Tyr | Met | Lys | Lys | Gly | Ser |
|    |     |     |     | 100 |     |     |     |     | 105 |     |     |     |
|    | Lys | Val | Met | Pro | Asn | Ile | Glu | Lys | Leu | Arg | Ser | Cys |
|    | 110 |     |     |     |     |     | 115 |     |     |     |     | 120 |
|    | Gly | Thr | His | Ser | Pro | Tyr | Met | Arg | Pro | Val | Tyr | Pro |
|    |     |     |     |     | 125 |     |     |     |     | 130 |     |     |
|    | Thr | Lys | Thr | Phe | Pro | Asn | Leu | Tyr | Thr | Leu | Ala | Thr |
|    |     |     | 135 |     |     |     |     | 140 |     |     |     |     |
| 5  | Gly | Leu | Tyr | Pro | Glu | Ser | His | Gly | Ile | Val | Gly | Asn |
|    | 145 |     |     |     |     | 150 |     |     |     |     | 155 |     |
|    | Ser | Met | Tyr | Asp | Pro | Val | Phe | Asp | Ala | Thr | Phe | His |
|    |     |     |     | 160 |     |     |     |     | 165 |     |     |     |
|    | Leu | Arg | Gly | Arg | Glu | Lys | Phe | Asn | His | Arg | Trp | Trp |
|    | 170 |     |     |     |     |     | 175 |     |     |     |     | 180 |
|    | Ala | Gly | Gln | Pro | Leu | Trp | Ile | Thr | Ala | Thr | Lys | Gln |
| 10 |     |     |     |     | 185 |     |     |     |     | 190 |     |     |
|    | Arg | Gly | Glu | Ser | Trp | Asn | Ile | Leu | Leu | Val | Cys | Cys |
|    |     |     | 195 |     |     |     |     | 200 |     |     |     |     |
|    | His | Pro | Ser | Arg | Ala | Glu | Ile | Leu | Thr | Ile | Leu | Gln |
|    | 205 |     |     |     |     | 210 |     |     |     |     | 215 |     |
|    | Trp | Leu | Thr | Leu | Pro | Asp | His | Glu | Arg | Pro | Ser | Val |
|    |     |     |     | 220 |     |     |     |     | 225 |     |     |     |
|    | Tyr | Ala | Phe | Tyr | Ser | Glu | Gln | Pro | Asp | Phe | Ser | Gly |
| 15 |     | 230 |     |     |     |     | 235 |     |     |     |     | 240 |
|    | His | Lys | His | Met | Pro | Phe | Gly | Pro | Glu | Met | Pro | Asn |
|    |     |     |     | 245 |     |     |     |     |     | 250 |     |     |
|    | Pro | Leu | Arg | Glu | Met | His | Lys | Ile | Val | Gly | Gln | Leu |
|    |     |     | 255 |     |     |     |     | 260 |     |     |     |     |
|    | Met | Asp | Gly | Leu | Lys | Gln | Leu | Lys | Leu | His | Arg | Cys |
|    | 265 |     |     |     |     | 270 |     |     |     |     | 275 |     |
| 20 | Val | Asn | Val | Ile | Phe | Val | Glu | Thr | Met | Asp | Gly | Arg |
|    |     |     |     | 280 |     |     |     |     | 285 |     |     |     |
|    | Cys | His | Met | Tyr | Arg | Thr | Glu | Phe | Leu | Ser | Asn | Tyr |
|    |     | 290 |     |     |     |     | 295 |     |     |     |     | 300 |
|    | Leu | Thr | Asn | Val | Asp | Asp | Ile | Thr | Leu | Val | Pro | Gly |
|    |     |     |     | 305 |     |     |     |     |     | 310 |     |     |
|    | Thr | Leu | Gly | Arg | Ile | Arg | Ser | Lys | Phe | Ser | Asn | Asn |
|    |     |     | 315 |     |     |     |     | 320 |     |     |     |     |
| 25 | Ala | Lys | Tyr | Asp | Pro | Lys | Ala | Ile | Ile | Ala | Asn | Leu |
|    | 325 |     |     |     |     | 330 |     |     |     |     | 335 |     |
|    | Thr | Cys | Lys | Lys | Pro | Asp | Gln | His | Phe | Lys | Pro | Tyr |
|    |     |     |     | 340 |     |     |     |     | 345 |     |     |     |
|    | Leu | Lys | Gln | His | Leu | Pro | Lys | Arg | Leu | His | Tyr | Ala |
|    |     | 350 |     |     |     |     | 355 |     |     |     |     | 360 |
|    | Asn | Asn | Arg | Arg | Ile | Glu | Asp | Ile | His | Leu | Leu | Val |

- 64 -

|    |     |     |     |     |     |     |     |     |     |     |     |     |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
|    | Tyr | Lys | Thr | Lys | Val | Pro | Pro | Phe | Glu | Asn | Ile | Glu |
|    |     |     |     |     | 425 |     |     |     |     | 430 |     |     |
|    | Leu | Tyr | Asn | Val | Met | Cys | Asp | Leu | Leu | Gly | Leu | Lys |
|    |     |     | 435 |     |     |     |     | 440 |     |     |     |     |
|    | Pro | Ala | Pro | Asn | Asn | Gly | Thr | His | Phe | Ser | Leu | Asn |
|    | 445 |     |     |     |     | 450 |     |     |     |     | 455 |     |
| 5  | His | Leu | Leu | Arg | Thr | Asn | Thr | Phe | Arg | Pro | Thr | Met |
|    |     |     |     | 460 |     |     |     |     | 465 |     |     |     |
|    | Pro | Glu | Glu | Val | Thr | Arg | Pro | Asn | Tyr | Pro | Gly | Ile |
|    | 470 |     |     |     |     |     | 475 |     |     |     |     | 480 |
|    | Met | Tyr | Leu | Gln | Ser | Asp | Phe | Asp | Leu | Gly | Cys | Thr |
|    |     |     |     |     | 485 |     |     |     |     | 490 |     |     |
|    | Cys | Asp | Asp | Lys | Val | Glu | Pro | Lys | Asn | Lys | Leu | Asp |
|    |     |     | 495 |     |     |     |     | 500 |     |     |     |     |
| 10 | Glu | Leu | Asn | Lys | Arg | Leu | His | Thr | Lys | Gly | Ser | Thr |
|    | 505 |     |     |     |     | 510 |     |     |     |     | 515 |     |
|    | Glu | Glu | Arg | His | Leu | Leu | Tyr | Gly | Asp | Arg | Pro | Ala |
|    |     |     |     | 520 |     |     |     |     | 525 |     |     |     |
|    | Val | Leu | Tyr | Arg | Thr | Arg | Tyr | Asp | Ile | Leu | Tyr | His |
|    | 530 |     |     |     |     |     | 535 |     |     |     |     | 540 |
|    | Thr | Asp | Phe | Glu | Ser | Gly | Tyr | Ser | Glu | Ile | Phe | Leu |
|    |     |     |     |     | 545 |     |     |     |     | 550 |     |     |
| 15 | Met | Pro | Leu | Trp | Thr | Ser | Tyr | Thr | Val | Ser | Lys | Gln |
|    |     |     | 555 |     |     |     |     | 560 |     |     |     |     |
|    | Ala | Glu | Val | Ser | Ser | Val | Pro | Asp | His | Leu | Thr | Ser |
|    | 565 |     |     |     |     | 570 |     |     |     |     | 575 |     |
|    | Cys | Val | Arg | Pro | Asp | Val | Arg | Val | Ser | Pro | Ser | Phe |
|    |     |     |     | 580 |     |     |     |     | 585 |     |     |     |
|    | Ser | Gln | Asn | Cys | Leu | Ala | Tyr | Lys | Asn | Asp | Lys | Gln |
|    | 590 |     |     |     |     |     | 595 |     |     |     |     | 600 |
| 20 | Met | Ser | Tyr | Gly | Gly | Leu | Gly | Pro | Pro | Tyr | Leu | Ser |
|    |     |     |     | 605 |     |     |     |     |     | 610 |     |     |
|    | Ser | Ser | Pro | Glu | Ala | Lys | Tyr | Asp | Ala | Phe | Leu | Val |
|    |     |     | 615 |     |     |     |     | 620 |     |     |     |     |
|    | Thr | Asn | Met | Val | Pro | Met | Tyr | Pro | Ala | Phe | Lys | Arg |
|    | 625 |     |     |     |     | 630 |     |     |     |     | 635 |     |
| 25 | Val | Trp | Asn | Tyr | Phe | Gln | Arg | Val | Leu | Val | Lys | Lys |
|    |     |     |     | 640 |     |     |     |     | 645 |     |     |     |
|    | Tyr | Ala | Ser | Glu | Arg | Asn | Gly | Val | Asn | Val | Ile | Ser |
|    | 650 |     |     |     |     |     | 655 |     |     |     |     | 660 |
|    | Gly | Pro | Ile | Phe | Asp | Tyr | Asp | Tyr | Asp | Gly | Leu | His |
|    |     |     |     | 665 |     |     |     |     |     | 670 |     |     |
|    | Asp | Thr | Glu | Asp | Lys | Ile | Lys | Gln | Tyr | Val | Glu | Gly |
|    |     |     | 675 |     |     |     |     | 680 |     |     |     |     |
| 30 | Ser | Ser | Ile | Pro | Val | Pro | Thr | His | Tyr | Tyr | Ser | Ile |
|    | 685 |     |     |     |     | 690 |     |     |     |     | 695 |     |
|    | Ile | Thr | Ser | Cys | Leu | Asp | Phe | Thr | Gln | Pro | Ala | Asp |
|    |     |     |     | 700 |     |     |     |     | 705 |     |     |     |
|    | Lys | Cys | Asp | Gly | Pro | Leu | Ser | Val | Ser | Ser | Phe | Ile |
|    | 710 |     |     |     |     |     | 715 |     |     |     |     | 720 |
|    | Leu | Pro | His | Arg | Pro | Asp | Asn | Glu | Glu | Ser | Cys | Asn |
|    |     |     |     |     | 725 |     |     |     |     |     | 730 |     |
| 35 |     |     |     |     |     |     |     |     |     |     |     |     |

- 65 -

° Ser Ser Glu Asp Glu Ser Lys Trp Val Glu Glu Leu  
                     735                    740  
 Met Lys Met His Thr Ala Arg Val Arg Asp Ile Glu  
 745                    750                    755  
 His Leu Thr Ser Leu Asp Phe Phe Arg Lys Thr Ser  
                     760                    765  
 Arg Ser Tyr Pro Glu Ile Leu Thr Leu Lys Thr Tyr  
 5       770                    775                    780  
 Leu His Thr Tyr Glu Ser Glu Ile  
                     785

## (2) INFORMATION FOR SEQ ID NO:37:

- 10 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 2712  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: Unknown
- (ii) MOLECULE TYPE: cDNA
- 15 (iii) HYPOTHETICAL: No
- (vi) ORIGINAL SOURCE:  
 (A) ORGANISM: Human  
 (B) STRAIN:  
 (C) INDIVIDUAL ISOLATE:  
 (D) DEVELOPMENTAL STAGE:  
 (E) HAPLOTYPE:  
 20 (F) TISSUE TYPE:  
 (G) CELL TYPE: teratocarcinoma  
 (H) CELL LINE: N-tera 2D1  
 (I) ORGANELLE:
- (ix) FEATURE:  
 (A) NAME/KEY:  
 25 (B) LOCATION:  
 (C) IDENTIFICATION METHOD:  
 (D) OTHER INFORMATION: N-tera 2D1 ATX DNA  
                     sequence

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

|    |            |            |            |            |     |
|----|------------|------------|------------|------------|-----|
|    | TGTGACAACT | TGTGTAAGAG | CTATACCACT | TGCTGCCATG | 40  |
| 30 | ACTTTGATGA | GCTGTGTTTG | AAGACAGCCC | GTGCGTGCGA | 80  |
|    | GTGTACTAAG | GACAGATGTG | GGAAGTCAG  | AAATGAAGAA | 120 |
|    | AATGCCTGTC | ACTGCTCAGA | GGAAGTCAG  | GCCAGGGGAG | 160 |
|    | ACTGCTGTAA | CAATTACCAA | GTGGTTTGCA | AAGGAGAGTC | 200 |
|    | GCATTGGGTT | GATGATGACT | GTGAGGAAAT | AAAGGCCGCA | 240 |
|    | GAATGCCTGC | AGGTTTGTTT | GCCCTCCATT | AATCATCTTC | 280 |
|    | TCCGTGGATG | GCTTCCGATG | ACATCATACA | TGAAGAAAGG | 320 |
|    | CAGCAAAGTC | ATGCCTAATA | TTGAAAAACT | AAGGTCTTGT | 360 |
| 35 | GGCACACACT | CTCCCTACAT | GAGGCCGGTG | TACCCAATA  | 400 |

- 66 -

|    |             |             |             |            |      |
|----|-------------|-------------|-------------|------------|------|
|    | AAACCTTTCC  | TAACTTATAC  | ACTTTGGCCA  | CTGGGCTATA | 440  |
|    | TCCAGAATCA  | CATGGAATTG  | TTGGCAATTC  | AATGTATGAT | 480  |
|    | CCTGTATTTG  | ATGCCACTTT  | TCATCTGCGA  | GGGCGAGAGA | 520  |
|    | AATTTAATCA  | TAGATGGTGG  | GGAGGTCAAC  | CGCTATGGAT | 560  |
|    | TACAGCCACC  | AAGCAAAGGG  | GTGAAAGCTG  | GAACATTCTT | 600  |
|    | TTGGTCTGTT  | GTCATCCCTC  | ACGAGCGGAG  | ATATTAACCA | 640  |
| 5  | TATTGCAGTG  | GCTCACCTG   | CCAGATCATG  | AGAGGCCTTC | 680  |
|    | GGTCTATGCC  | TTCTATTCTG  | AGCAACCTGA  | TTTCTCTGGA | 720  |
|    | CACAAACATA  | TGCCTTTTCGG | CCCTGAGATG  | ACAAATCCTC | 760  |
|    | TGAGGGGAAAT | GCACAAAATT  | GTGGGGCAAT  | TAATGGATGG | 800  |
|    | ACTGAAACAA  | CTAAAACTGC  | ATCGGTGTGT  | CAACGTCATC | 840  |
|    | TTTGTGCGAGA | CCATGGATGG  | AAGATGTAC   | ATGTATAGAA | 880  |
|    | CTGAGTTCTT  | GAGTAATTAC  | CTAACTAATG  | TGGATGATAT | 920  |
|    | TACTTTAGTG  | CCTGGAACCTC | TAGGAAGAAT  | TCGATCCAAA | 960  |
| 10 | TTTAGCAACA  | ATGCTAAATA  | TCACCCCAAA  | GCCATTATTG | 1000 |
|    | CCAATCTCAC  | GTGTAAAAAA  | CCAGATCAGC  | ACTTTAAGCC | 1040 |
|    | TTACTTGAAA  | CAGCACCTTC  | CCAAACGTTT  | GCACTATGCC | 1080 |
|    | AACAACAGAA  | GAATTGAGGA  | TATCCATTTA  | TTGGTGGAAC | 1120 |
|    | GCAGATGGCA  | TGTTGCAAGG  | AAACCTTTGG  | ATGTTTATAA | 1160 |
|    | GAAACCATCA  | GGAAATGCTT  | TTTCCAGGGA  | GACCACGGCA | 1200 |
|    | TTTGATAACA  | AGGTCAACAG  | CATGCAGACT  | GTTTTTGTAG | 1240 |
| 15 | GTTATGGCCC  | AACATTTAAG  | TACAAGACTA  | AAGTDCCTCC | 1280 |
|    | ATTTGAAAAC  | ATTGAACTTT  | AAAATGTTAT  | GTGTGATCTC | 1320 |
|    | CTGGGATTGA  | AGCCAGCTCC  | TAATAATGGG  | ACCCATGGAA | 1360 |
|    | GTTTGAATCA  | TCTCCTGCGC  | ACTAATACCT  | TCAGGCCAAC | 1400 |
|    | CATGCCAGAG  | GAAGTTACCA  | GACCCTATTA  | TCCAGGGATT | 1440 |
|    | ATGTACCTTC  | AGTCTGATTT  | TGACCTGGGC  | TGCACTTGTG | 1480 |
|    | ATGATAAGGT  | AGAGCCAAAG  | AACAAGTTGG  | ATGAACTCAA | 1520 |
|    | CAACCGGCTT  | CATACAAAAG  | GGTCTACAGA  | AGAGAGACAC | 1560 |
| 20 | CTCCTCTATG  | GGGATCGACC  | TGCAGTGCTT  | TATCGGACTA | 1600 |
|    | GATATGATAT  | CTTATATCAC  | ACTGACTTTG  | AAAGTGGTTA | 1640 |
|    | TAGTGAAATA  | TTCTTAATGC  | CACTCTGGAC  | ATCATATACT | 1680 |
|    | GTTTCCAAAC  | AGGCTGAGGT  | TTCCAGCGTT  | CCTGACCATC | 1720 |
|    | TGACCAGTTG  | CGTCCGGCCT  | GATGTCCGTG  | TTTCTCCGAG | 1760 |
|    | TTTCAGTCAG  | AACTGTTTGG  | CCTACAAAAA  | TGATAAGCAG | 1800 |
|    | ATGTCCTACG  | GATTCCTCTT  | TCCTCCTTAT  | CTGAGCTCTT | 1840 |
| 25 | CACCAGAGGC  | TAAATATGAT  | GCATTCTTTG  | TAACCAATAT | 1880 |
|    | GGTTCCAATG  | TATCCTGCTT  | TCAAACGGGT  | CTGGAATTAT | 1920 |
|    | TTCCAAAGGG  | TATTGGTGAA  | GAAATATGCT  | TCGGAAAGAA | 1960 |
|    | ATGGAGTTAA  | CGTGATAAGT  | GGACCAATCT  | TCGACTATGA | 2000 |
|    | CTATGATGGC  | TTACATGACA  | CAGAAGACAA  | AATAAAACAG | 2040 |
|    | TACGTGGAAG  | GCAGTTCCAT  | TCCTGTTCCA  | ACTCACTACT | 2080 |
|    | ACAGCATCAT  | CACCAGCTGT  | CTGGATTTCA  | CTCAGCCTGC | 2120 |
|    | CGACAAGTGT  | GACGGCCCTC  | TCTCTGTGTC  | CTCCTTCATC | 2160 |
| 30 | CTGCCTCACC  | GGCCTGACAA  | CGAGGAGAGC  | TGCAATAGCT | 2200 |
|    | CAGAGGACGA  | ATCAAAATGG  | GTAAGAAGAAC | TCATGAAGAT | 2240 |
|    | GCACACAGCT  | AGGGTGCGTG  | ACATTGAACA  | TCTCACCAGC | 2280 |
|    | CTGGACTTCT  | TCCGAAAGAC  | CAGCCGCAGC  | TACCCAGAAA | 2320 |
|    | TCCTGACACT  | CAAGACATAC  | CTGCATACAT  | ATGAGAGCGA | 2360 |
|    | GATTTAACTT  | TCTGAGCATC  | TGCAGTACAG  | TCTTATCAAC | 2400 |
|    | TGGTTGTATA  | TTTTTATATT  | GTTTTTGTAT  | TTATTAATTT | 2440 |
| 35 | GAAACCAGGA  | CATTAAAAAT  | GTTAGTATTT  | TAATCCTGTA | 2480 |
|    | CCAAATCTGA  | CATATTATGC  | CTGAATGACT  | CCACTGTTTT | 2520 |



- 67 -

TCTCTAATGC TTGATTTAGG TAGCCTTGTG TTCTGAGTAG 2560  
 AGCTTGTAAT AAATACTGCA GCTTGAGTTT TTAGTGGAAG 2600  
 CTTCTAAATG GTGCTGCAGA TTTGATATTT GCATTGAGGA 2640  
 AATATTAATT TTCCAATGCA CAGTTGCCAC ATTTAGTCCT 2680  
 GTACTGTATG GAAACACTGA TTTTGTAAG TT 2712

5 (2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 979  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: Unknown

10 (ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: No

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Human  
 (B) STRAIN:  
 (C) INDIVIDUAL ISOLATE:  
 (D) DEVELOPMENTAL STAGE:  
 (E) HAPLOTYPE:  
 (F) TISSUE TYPE: Liver  
 (G) CELL TYPE:  
 (H) CELL LINE:  
 (I) ORGANELLE:

20 (ix) FEATURE:

- (A) NAME/KEY:  
 (B) LOCATION:  
 (C) IDENTIFICATION METHOD:  
 (D) OTHER INFORMATION: putative autotaxin  
 protein sequence from human liver

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

Met Ala Arg Arg Ser Ser Phe Gln Ser Cys Gln Asp  
 1 5 10  
 Ile Ser Leu Phe Thr Phe Ala Val Gly Val Asn Ile  
 15 20  
 Cys Leu Gly Phe Thr Ala His Arg Ile Lys Arg Ala  
 25 30 35  
 Glu Gly Trp Glu Glu Gly Pro Pro Thr Val Leu Ser  
 40 45  
 Asp Ser Pro Trp Thr Asn Ile Ser Gly Ser Cys Lys  
 50 55 60  
 Gly Arg Cys Phe Glu Leu Gln Glu Ala Gly Pro Pro  
 65 70  
 Asp Cys Arg Cys Asp Asn Leu Cys Lys Ser Tyr Thr  
 75 80

- 68 -

Ser Cys Cys His Asp Phe Asp Glu Leu Cys Leu Lys  
 85 90 95  
 Thr Ala Arg Ala Trp Glu Cys Thr Lys Asp Arg Cys  
 100 105  
 Gly Glu Val Arg Asn Glu Glu Asn Ala Cys His Cys  
 110 115 120  
 5 Ser Glu Asp Cys Leu Ala Arg Gly Asp Cys Cys Thr  
 125 130  
 Asn Tyr Gln Val Val Cys Lys Gly Glu Ser His Trp  
 135 140  
 Val Asp Asp Asp Cys Glu Glu Ile Lys Ala Ala Glu  
 145 150 155  
 Cys Leu Gln Val Cys Ser Pro Ser Ile Asn His Leu  
 160 165  
 10 Leu Arg Gly Trp Leu Pro Met Thr Ser Tyr Met Lys  
 170 175 180  
 Lys Gly Ser Lys Val Met Pro Asn Ile Glu Lys Leu  
 185 190  
 Arg Ser Cys Gly Thr His Ser Pro Tyr Met Arg Pro  
 195 200  
 Val Tyr Pro Thr Lys Thr Phe Pro Asn Leu Tyr Thr  
 205 210 215  
 15 Leu Ala Thr Gly Leu Tyr Pro Glu Ser His Gly Ile  
 220 225  
 Val Gly Asn Ser Met Tyr Asp Pro Val Phe Asp Ala  
 230 235 240  
 Thr Phe His Leu Arg Gly Arg Glu Lys Phe Asn His  
 245 250  
 Arg Trp Trp Gly Gly Gln Pro Leu Trp Ile Thr Ala  
 255 260  
 20 Thr Lys Gln Arg Gly Glu Ser Trp Asn Ile Leu Leu  
 265 270 275  
 Val Cys Cys His Pro Ser Arg Ala Glu Ile Leu Thr  
 280 285  
 Ile Leu Gln Trp Leu Thr Leu Pro Asp His Glu Arg  
 290 295 300  
 25 Pro Ser Val Tyr Ala Phe Tyr Ser Glu Gln Pro Asp  
 305 310  
 Phe Ser Gly His Lys His Met Pro Phe Gly Pro Glu  
 315 320  
 Met Thr Asn Pro Leu Arg Glu Met His Lys Ile Val  
 325 330 335  
 Gly Gln Leu Met Asp Gly Leu Lys Gln Leu Lys Leu  
 340 345  
 30 His Arg Cys Val Asn Val Ile Phe Val Glu Thr Met  
 350 355 360  
 Asp Gly Arg Cys His Met Tyr Arg Thr Glu Phe Leu  
 365 370  
 Ser Asn Tyr Leu Thr Asn Val Asp Asp Ile Thr Leu  
 375 380  
 Val Pro Gly Thr Leu Gly Arg Ile Arg Ser Lys Phe  
 385 390 395  
 35

- 69 -

° Ser Asn Asn Ala Lys Tyr Asp Pro Lys Ala Ile Ile  
                     400                    405  
 Ala Asn Leu Thr Cys Lys Lys Pro Asp Gln His Phe  
           410                    415                    420  
 Lys Pro Tyr Leu Lys Gln His Leu Pro Lys Arg Leu  
                     425                    430  
 His Tyr Ala Asn Asn Arg Arg Ile Glu Asp Ile His  
                     435                    440  
 5 Leu Leu Val Glu Arg Arg Trp His Val Ala Arg Lys  
       445                    450                    455  
 Pro Leu Asp Val Tyr Lys Lys Pro Ser Gly Asn Ala  
                     460                    465  
 Phe Ser Arg Glu Thr Thr Ala Phe Asp Asn Lys Val  
           470                    475                    480  
 Asn Ser Met Gln Thr Val Phe Val Gly Tyr Gly Pro  
                     485                    490  
 10 Thr Phe Lys Tyr Lys Thr Lys Val Pro Pro Phe Glu  
                     495                    500  
 Asn Ile Glu Leu Tyr Asn Val Met Cys Asp Leu Leu  
       505                    510                    515  
 Gly Leu Lys Pro Ala Pro Asn Asn Gly Thr His Gly  
                     520                    525  
 15 Ser Leu Asn His Leu Leu Arg Thr Asn Thr Phe Arg  
       530                    535                    540  
 Pro Thr Met Pro Glu Glu Val Thr Arg Pro Asn Tyr  
                     545                    550  
 Pro Gly Ile Met Tyr Leu Gln Ser Asp Phe Asp Leu  
                     555                    560  
 Gly Cys Thr Cys Asp Asp Lys Val Glu Pro Lys Asn  
       565                    570                    575  
 20 Lys Leu Asp Glu Leu Asn Lys Arg Leu His Thr Lys  
                     580                    585  
 Gly Ser Thr Glu Glu Arg His Leu Leu Tyr Gly Asp  
           590                    595                    600  
 Arg Pro Ala Val Leu Tyr Arg Thr Arg Tyr Asp Ile  
                     605                    610  
 Leu Tyr His Thr Asp Phe Glu Ser Gly Tyr Ser Glu  
                     615                    620  
 25 Ile Phe Leu Met Pro Leu Trp Thr Ser Tyr Thr Val  
       625                    630                    635  
 Ser Lys Gln Ala Glu Val Ser Ser Val Pro Asp His  
                     640                    645  
 Leu Thr Ser Cys Val Arg Pro Asp Val Arg Val Ser  
           650                    655                    660  
 Pro Ser Phe Ser Gln Asn Cys Leu Ala Tyr Lys Asn  
                     665                    670  
 30 Asp Lys Gln Met Ser Tyr Gly Phe Leu Phe Pro Pro  
                     675                    680  
 Tyr Leu Ser Ser Ser Pro Glu Ala Lys Tyr Asp Ala  
       685                    690                    695  
 Phe Leu Val Thr Asn Met Val Pro Met Tyr Pro Ala  
                     700                    705  
 35 Phe Lys Arg Val Trp Asn Tyr Phe Gln Arg Val Leu  
       710                    715                    720

- 70 -



|    |            |     |            |            |            |            |            |            |            |            |            |            |
|----|------------|-----|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
|    | Val        | Lys | Lys        | Tyr        | Ala<br>725 | Ser        | Glu        | Arg        | Asn<br>730 | Gly        | Val        | Asn        |
|    | Val        | Ile | Ser<br>735 | Gly        | Pro        | Ile        | Phe        | Asp<br>740 | Tyr        | Asp        | Tyr        | Asp        |
|    | Gly<br>745 | Leu | His        | Asp        | Thr        | Glu<br>750 | Asp        | Lys        | Ile        | Lys        | Gln        | Tyr<br>755 |
| 5  | Val        | Glu | Gly        | Ser<br>760 | Ser        | Ile        | Pro        | Val        | Pro<br>765 | Thr        | His        | Tyr        |
|    | Tyr<br>770 | Ser | Ile        | Ile        | Thr        | Ser        | Cys<br>775 | Leu        | Asp        | Phe        | Thr        | Gln<br>780 |
|    | Pro        | Ala | Asp        | Lys<br>785 | Cys        | Asp        | Gly        | Pro        | Leu        | Ser<br>790 | Val        | Ser        |
|    | Ser        | Phe | Ile<br>795 | Leu        | Pro        | His        | Arg        | Pro<br>800 | Asp        | Asn        | Glu        | Glu        |
| 10 | Ser<br>805 | Cys | Asn        | Ser        | Ser        | Glu<br>810 | Asp        | Glu        | Ser        | Lys        | Trp<br>815 | Val        |
|    | Glu        | Glu | Leu        | Met<br>820 | Lys        | Met        | His        | Thr        | Ala<br>825 | Arg        | Val        | Arg        |
|    | Asp<br>830 | Ile | Glu        | His        | Leu        | Thr        | Ser<br>835 | Leu        | Asp        | Phe        | Phe        | Arg<br>840 |
|    | Lys        | Thr | Ser        | Arg        | Ser<br>845 | Tyr        | Pro        | Glu        | Ile        | Leu        | Thr        | Leu        |
| 15 | Lys        | Thr | Tyr<br>855 | Leu        | His        | Thr        | Tyr        | Glu<br>860 | Ser        | Glu        | Ile        | Xaa        |
|    | Leu<br>865 | Ser | Glu        | His        | Leu        | Gln<br>870 | Tyr        | Ser        | Leu        | Ile        | Asn<br>875 | Trp        |
|    | Leu        | Tyr | Ile<br>880 | Phe        | Ile        | Leu        | Phe        | Leu<br>885 | Tyr        | Leu        | Leu        | Ile        |
|    | Xaa<br>890 | Asn | Gln        | Asp        | Ile        | Lys        | Asn<br>895 | Val        | Ser        | Ile        | Leu        | Ile<br>900 |
| 20 | Leu        | Tyr | Gln        | Ile        | Xaa<br>905 | His        | Ile        | Met        | Pro        | Glu<br>910 | Xaa        | Leu        |
|    | His        | Cys | Phe<br>915 | Ser        | Leu        | Met        | Leu        | Asp<br>920 | Leu        | Gly        | Ser        | Leu        |
|    | Val<br>925 | Phe | Xaa        | Val        | Glu        | Leu<br>930 | Val        | Ile        | Asn        | Thr        | Ala        | Ala<br>935 |
| 25 | Xaa        | Val | Phe        | Ser<br>940 | Gly        | Ser        | Phe        | Xaa<br>945 | Met        | Val        | Leu        | Gln        |
|    | Ile<br>950 | Xaa | Tyr        | Leu        | His        | Xaa        | Gly<br>955 | Asn        | Ile        | Asn        | Phe        | Pro<br>960 |
|    | Met        | His | Ser        | Cys<br>965 | His        | Ile        | Xaa        | Ser        | Cys        | Thr<br>970 | Val        | Trp        |
|    | Lys        | His | Xaa<br>975 | Phe        | Cys        | Lys        | Val        |            |            |            |            |            |

30

(2) INFORMATION FOR SEQ ID NO:39:

```
(i) SEQUENCE CHARACTERISTICS:
    (A) LENGTH:8
    (B) TYPE: amino acids
    (C) STRANDEDNESS: single
```

35

- 71 -

- ° (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE:  
(A) DESCRIPTION: peptide
- (iii) HYPOTHETICAL: No
- 5 (ix) FEATURE:  
(A) NAME/KEY: ATX-204  
(B) LOCATION:  
(C) IDENTIFICATION METHOD:  
(D) OTHER INFORMATION:
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:
- 10 Met His Thr Ala Arg Val Arg Asp  
5
- (2) INFORMATION FOR SEQ ID NO:40:
- 15 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 8  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: No
- 20 (ix) FEATURE:  
(A) NAME/KEY: ATX-205  
(B) LOCATION:  
(C) IDENTIFICATION METHOD:  
(D) OTHER INFORMATION:
- 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:
- Phe Ser Asn Asn Ala Lys Tyr Asp  
5
- (2) INFORMATION FOR SEQ ID NO:41:
- 30 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 7  
(B) TYPE: amino acids  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE:  
(A) DESCRIPTION: Peptide
- 35

- 72 -

°

(iii) HYPOTHETICAL: No

(ix) FEATURE:

- (A) NAME/KEY: ATX-209
- (B) LOCATION:
- (C) IDENTIFICATION METHOD:
- (D) OTHER INFORMATION:

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

Val Met Pro Asn Ile Glu Lys  
5

10 (2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8
- (B) TYPE: amino acids
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE:

- (A) DESCRIPTION: Peptide

(iii) HYPOTHETICAL: No

(ix) FEATURE:

- (A) NAME/KEY: ATX-210
- (B) LOCATION:
- (C) IDENTIFICATION METHOD:
- (D) OTHER INFORMATION:

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

25

Thr Ala Arg Gly Trp Glu Cys Thr  
5

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE:

- (A) DESCRIPTION: Peptide

35

(iii) HYPOTHETICAL: No

- 73 -

°

- (ix) FEATURE:  
(A) NAME/KEY: ATX-212  
(B) LOCATION:  
(C) IDENTIFICATION METHOD:  
(D) OTHER INFORMATION:

5

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

Xaa Asp Ser Pro Trp Thr Xaa Ile Ser Gly Ser  
5 10

10

- (2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 11  
(B) TYPE: amino acids  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

15

- (ii) MOLECULE TYPE:  
(A) DESCRIPTION: Peptide

- (iii) HYPOTHETICAL: No

20

- (ix) FEATURE:  
(A) NAME/KEY: ATX-214  
(B) LOCATION:  
(C) IDENTIFICATION METHOD:  
(D) OTHER INFORMATION:

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

25

Leu Arg Ser Cys Gly Thr His Ser Pro Tyr Met  
5 10

- (2) INFORMATION FOR SEQ ID NO:45:

30

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 8  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE:  
(A) DESCRIPTION: Peptide

- (iii) HYPOTHETICAL: No

35

- (ix) FEATURE:

- 74 -

- (A) NAME/KEY: ATX-215/34A
- (B) LOCATION:
- (C) IDENTIFICATION METHOD:
- (D) OTHER INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

5 Thr Tyr Leu His Thr Tyr Glu Ser  
5

(2) INFORMATION FOR SEQ ID NO:46:

- 10 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 13  
(B) TYPE: amino acids  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE:  
(A) DESCRIPTION: Peptide

15 (iii) HYPOTHETICAL: No

- (ix) FEATURE:  
(A) NAME/KEY:  
(B) LOCATION:  
(C) IDENTIFICATION METHOD:  
(D) OTHER INFORMATION:

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

Ala Ile Ile Ala Asn Leu Thr Cys Lys Lys Pro Asp Gln  
5 10

25 (2) INFORMATION FOR SEQ ID NO:47:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 8  
(B) TYPE: amino acids  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE:  
(A) DESCRIPTION: Peptide

(iii) HYPOTHETICAL: No

- (ix) FEATURE:  
(A) NAME/KEY: ATX-216  
(B) LOCATION:  
(C) IDENTIFICATION METHOD:

35



- 75 -

## (D) OTHER INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

Ile Val Gly Gln Leu Met Asp Gly  
5

5

(2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9
- (B) TYPE: amino acids
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE:

- (A) DESCRIPTION: Peptide

(iii) HYPOTHETICAL: No

(ix) FEATURE:

- (A) NAME/KEY: ATX-218/44
- (B) LOCATION:
- (C) IDENTIFICATION METHOD:
- (D) OTHER INFORMATION:

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

20 Thr Ser Arg Ser Tyr Pro Glu Ile Leu  
5

(2) INFORMATION FOR SEQ ID NO:49:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9
- (B) TYPE: amino acids
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE:

- (A) DESCRIPTION: Peptide

30

(iii) HYPOTHETICAL: No

(ix) FEATURE:

- (A) NAME/KEY: ATX-223B/24
- (B) LOCATION:
- (C) IDENTIFICATION METHOD:
- (D) OTHER INFORMATION:

35

- 76 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:  
Gln Ala Glu Val Ser Ser Val Pro Asp  
5

5 (2) INFORMATION FOR SEQ ID NO:50:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 14  
(B) TYPE: amino acids  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
10 (ii) MOLECULE TYPE:  
(A) DESCRIPTION: Peptide  
(iii) HYPOTHETICAL: No  
(ix) FEATURE:  
(A) NAME/KEY: ATX-224  
15 (B) LOCATION:  
(C) IDENTIFICATION METHOD:  
(D) OTHER INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:  
Arg Cys Phe Glu Leu Gln Glu Ala Gly Pro Pro Asp Asp Cys  
5 10  
20

(2) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 12  
(B) TYPE: amino acid  
25 (C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(ii) MOLECULE TYPE:  
(A) DESCRIPTION: Peptide  
(iii) HYPOTHETICAL: No  
30 (ix) FEATURE:  
(A) NAME/KEY: ATX-229  
(B) LOCATION:  
(C) IDENTIFICATION METHOD:  
(D) OTHER INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:  
35

- 77 -

° Ser Tyr Thr Ser Cys Cys His Asp Phe Asp Glu Leu  
5 10

## (2) INFORMATION FOR SEQ ID NO:52:

- 5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 16  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE:  
(A) DESCRIPTION: Peptide
- 10 (iii) HYPOTHETICAL: No
- (ix) FEATURE:  
(A) NAME/KEY: ATX-224/53  
(B) LOCATION:  
(C) IDENTIFICATION METHOD:  
(D) OTHER INFORMATION:
- 15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:
- Gln Met Ser Tyr Gly Phe Leu Phe Pro Pro Tyr Leu  
1 5 10  
Ser Ser Ser Pro  
15

20

## (2) INFORMATION FOR SEQ ID NO:53:

- 25 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 117  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: Unknown
- (ii) MOLECULE TYPE:  
(A) DESCRIPTION: cDNA
- (iii) HYPOTHETICAL: No
- (iv) ANTI-SENSE:
- 30 (vi) ORIGINAL SOURCE:  
(A) ORGANISM: Human  
(B) STRAIN:  
(C) INDIVIDUAL ISOLATE:  
(D) DEVELOPMENTAL STAGE:  
(E) HAPLOTYPE:  
(F) TISSUE TYPE: Liver  
35 (G) CELL TYPE:

- 78 -

(H) CELL LINE:

(I) ORGANELLE:

(ix) FEATURE:

(A) NAME/KEY:

(B) LOCATION:

5

(C) IDENTIFICATION METHOD:

(D) OTHER INFORMATION: 5' end of human liver  
ATX gene

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

|    |   |     |
|----|---|-----|
|    | ATGGCAAGGA GGAGCTCGTT CCAGTCGTGT CAAGATATAT | 40  |
|    | CCCTGTTTAC TTTTGCCGTT GGAGTCAATA TCTGCTTAGG | 80  |
| 10 | ATTCAGTCA CATCGAATTA AGAGAGCAGA AGGATGG     | 117 |

(2) INFORMATION FOR SEQ ID NO:54:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 39

(B) TYPE: amino acids

15

(C) STRANDEDNESS: single

(D) TOPOLOGY: Unknown

(ii) MOLECULE TYPE:

(A) DESCRIPTION: Peptide

(iii) HYPOTHETICAL: No

20

(v) FRAGMENT TYPE: N-terminal fragment

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Human

(B) STRAIN:

(C) INDIVIDUAL ISOLATE:

(D) DEVELOPMENTAL STAGE:

25

(E) HAPLOTYPE:

(F) TISSUE TYPE: Liver

(G) CELL TYPE:

(H) CELL LINE:

(I) ORGANELLE:

(ix) FEATURE:

(A) NAME/KEY:

30

(B) LOCATION:

(C) IDENTIFICATION METHOD:

(D) OTHER INFORMATION: N-terminal region  
including transmembrane domain of liver  
ATX protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

35

- 79 -

Met Ala Arg Arg Ser Ser Phe Gln Ser Cys Gln Asp  
 1 5 10  
 Ile Ser Leu Phe Thr Phe Ala Val Gly Val Asn Ile  
 15 20  
 Cys Leu Gly Phe Thr Ala His Arg Ile Lys Arg Ala  
 25 30 35  
 Glu Gly Trp

5

## (2) INFORMATION FOR SEQ ID NO:55:

## (i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 21  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: No

15

## (iv) ANTI-SENSE: Yes

## (ix) FEATURE:

- (A) NAME/KEY:  
 (B) LOCATION:  
 (C) IDENTIFICATION METHOD:

20

- (D) OTHER INFORMATION: Primer from 5' end of  
 4C11

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

GCTCAGATAA GGAGGAAAGA G

21

## 25 (2) INFORMATION FOR SEQ ID NO:56:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

30

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: No

## (iv) ANTI-SENSE: Yes

## (ix) FEATURE:

35

- (A) NAME/KEY:  
 (B) LOCATION:

- 80 -

- (C) IDENTIFICATION METHOD:
- (D) OTHER INFORMATION: Nested primers from 4C11

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

5 GAATCCGTAG GACATCTGCT T 21

(2) INFORMATION FOR SEQ ID NO:57:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: No
- (iv) ANTI-SENSE: Yes
- (ix) FEATURE:
  - (A) NAME/KEY:
  - (B) LOCATION:
  - (C) IDENTIFICATION METHOD:
  - (D) OTHER INFORMATION: Nested primers from 4C11

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:  
TGTAGGCCAA ACAGTTCTGA C 21

(2) INFORMATION FOR SEQ ID NO:58:

- 25 (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 25
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- 30 (iii) HYPOTHETICAL: No
- (iv) ANTI-SENSE: No
- (ix) FEATURE:
  - (A) NAME/KEY:
  - (B) LOCATION:
  - 35 (C) IDENTIFICATION METHOD:

- 81 -

(D) OTHER INFORMATION: Nested sense primer  
deduced from ATX-101, wherein N is  
inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

5 AAYTCNATGC ARACNGTNTT YGTNG

25

(2) INFORMATION FOR SEQ ID NO:59:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 26  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: No

15 (iv) ANTI-SENSE: No

(ix) FEATURE:

(A) NAME/KEY:  
(B) LOCATION:  
(C) IDENTIFICATION METHOD:  
(D) OTHER INFORMATION: Nested primer of ATX  
-101, wherein N is inosine

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

TTYGTNGGNT AYGGNCCNAC NTTYAA

26

(2) INFORMATION FOR SEQ ID NO:60:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

30

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(ix) FEATURE:

35 (A) NAME/KEY:  
(B) LOCATION:  
(C) IDENTIFICATION METHOD:

- 82 -

(D) OTHER INFORMATION: Nested primer deduced from ATX-103, wherein N is inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

AAAYTAYCTNA CNAAYGTNGA YGAYAT

26

5

(2) INFORMATION FOR SEQ ID NO:61:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

15

(ix) FEATURE:

- (A) NAME/KEY:
- (B) LOCATION:
- (C) IDENTIFICATION METHOD:
- (D) OTHER INFORMATION: Nested primer deduced from ATX-103, wherein N is inosine

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

GAYGAYATNA CNCTNGTNCC NGGNAC

26

(2) INFORMATION FOR SEQ ID NO:62:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: No

30

(iv) ANTI-SENSE: No

(ix) FEATURE:

- (A) NAME/KEY:
- (B) LOCATION:
- (C) IDENTIFICATION METHOD:

35



- 83 -

(D) OTHER INFORMATION: Nested primer deduced  
from ATX-103, wherein N is inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

TGYTTYGARY TNCARGARGC NGGNCCNCC

29

5

(2) INFORMATION FOR SEQ ID NO:63:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

15

GCTGTCTTCA AACACAGC

18

(2) INFORMATION FOR SEQ ID NO:64:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21

(B) TYPE: nucleic acid

20

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: No

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

CTGGTGGCTG TAATCCATAG C

21

(2) INFORMATION FOR SEQ ID NO:65:

(i) SEQUENCE CHARACTERISTICS:

30

(A) LENGTH: 21

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

35

(iii) HYPOTHETICAL: No

- 84 -

- (iv) ANTI-SENSE: No
- (ix) FEATURE:
- (A) NAME/KEY:
  - (B) LOCATION:
  - (C) IDENTIFICATION METHOD:
  - (D) OTHER INFORMATION: Primer for 5' end of N-tera 2D1 sequence

5

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

CGTGAAGGCA AAGAGAACAC G

21

- 10 (2) INFORMATION FOR SEQ ID NO:66:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 3104
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: Unknown

15

- (ii) MOLECULE TYPE: cDNA

- (iii) HYPOTHETICAL: No

- (ix) FEATURE:
- (A) NAME/KEY: N-tera 2D1 ATX cDNA
  - (B) LOCATION:
  - (C) IDENTIFICATION METHOD:
  - (D) OTHER INFORMATION:

20

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

|    |             |            |            |            |            |     |
|----|-------------|------------|------------|------------|------------|-----|
|    | AGTGC       | ACTCC      | GTGAAGGCAA | AGAGAACACG | CTGCAAAAGG | 40  |
|    | CTTTCCAATA  | ATCCTCGACA | TGGCAAGGAG | GAGCTCGTTC |            | 80  |
| 25 | CAGTCGTGTC  | AGATAATATC | CCTGTTC    | ACTT       | TTTGCCGTTG | 120 |
|    | GAGTCAATAT  | CTGCTTAGGA | TTC        | ACTGCAC    | ATCGAATTAA | 160 |
|    | GAGAGCAGAA  | GGATGGGAGG | AAGGTCCTCC | TACAGTGCTA |            | 200 |
|    | TCAGACTCCC  | CCTGGACCAA | CATCTCCGGA | TCTTGCAAGG |            | 240 |
|    | GCAGGTGCTT  | TGAACTTCAA | GAGGCTGGAC | CTCCTGATTG |            | 280 |
|    | TCGCTGTGAC  | AACTTGTGTA | AGAGCTATAC | CAGTTGCTGC |            | 320 |
|    | CATGACTTTG  | ATGAGCTGTG | TTTGAAGACA | GCCCGTGCGT |            | 360 |
| 30 | GGGAGTGTAC  | TAAGGACAGA | TGTGGAGAAG | TCAGAAATGA |            | 400 |
|    | AGAAAATGCC  | TGTC       | ACTGCT     | CAGAGGACTG | CTTGCCAGG  | 440 |
|    | GGAGACTGCT  | GTACCAATTA | CCAAGTGGTT | TGCAAAGGAG |            | 480 |
|    | AGTCGCATTG  | GGTTGATGAT | GA         | CTGTGAGG   | AAATAAAGGC | 520 |
|    | CGCAGAAATGC | CCTGCAGGGT | TTGTT      | CGCCC      | TCCATTAATC | 560 |
|    | ATCTTCTCCG  | TGGATGGCTT | CCGTGCATCA | TACATGAAGA |            | 600 |
|    | AAGGCAGCAA  | AGTCATGCCT | AATATTGAAA | AACTAAGGTC |            | 640 |
|    | TTGTGGCACA  | CACTCGCCCC | ACATGAGGCC | GGTGTACCCA |            | 680 |
| 35 | ACTAAACCT   | TTCCTAACTT | ATACACTTTG | GCCACTGGGC |            | 720 |

- 85 -

|    |             |             |            |            |      |
|----|-------------|-------------|------------|------------|------|
|    | TATATCCAGA  | ATCACATGGA  | ATTGTTGGCA | ATTCAATGTA | 760  |
|    | TGATCCTGTA  | TTTGATGCCA  | CTTTTCATCT | GCGAGGGCGA | 800  |
|    | GAGAAATTTA  | ATCATAGATG  | GTGGGGAGGT | CAACCGCTAT | 840  |
|    | GGATTACAGC  | CACCAAGCAA  | AGGGGTGAAA | GCTGGAACAT | 880  |
|    | TCTTTTGGTC  | TGTTGTCATC  | CCTCACGAGC | GGAGATATTA | 920  |
|    | ACCATATTGC  | AGTGGCTCAC  | CCTGCCAGAT | CATGAGAGGC | 960  |
| 5  | TTCGGTCTAT  | GCCTTCTATT  | CTGAGCAACC | TGATTTCTCT | 1000 |
|    | GGACACAAAT  | ATGCCTTTTCG | GCCCTGAGAT | GACAAATCCT | 1040 |
|    | CTGAGGGAAA  | TCGACAAAAT  | TGTGGGGCAA | TTAATGGATG | 1080 |
|    | GACTGAAACA  | ACTAAAACCTG | CATCGGTGTG | TCAACGTCAT | 1120 |
|    | CTTTGTGCGA  | GACCATGGAA  | TGGAAGATGT | CACATGTGAT | 1160 |
|    | AGAACTGAGT  | TCTTGAGTAA  | TTACCTAACT | AATGTGGATG | 1200 |
|    | ATATTACTTT  | AGTGCCTGGA  | ACTCTAGGAA | TTCGATCCAA | 1240 |
|    | ATTTAGCAAC  | AATGCTAAAT  | ATGACCCCAA | AGCCATTATT | 1280 |
| 10 | GCCAATCTCA  | CGTGTA AAAA | ACCAGATCAG | CACTTTAAGC | 1320 |
|    | CTTACTTGAA  | ACAGCACCTT  | CCCAAACGTT | TGCACTATGC | 1360 |
|    | CAACAACAGA  | AGAATTGAGG  | ATATCCATTT | ATTGGTGGAA | 1400 |
|    | CGCAGATGGC  | ATGTTGCAAG  | GAAACCTTTG | GATGTTTATA | 1440 |
|    | AGAAACCATC  | AGGAAAATGC  | TTTTTCCAGG | GAGACCACGG | 1480 |
|    | ATTTGATAAC  | AAGGTCAACA  | GCATGCAGAC | TGTTTTTGTA | 1520 |
|    | GGTTATGGCC  | CAACATTTAA  | GTACAAGACT | AAAGTGCCTC | 1560 |
| 15 | CATTTGAAAA  | CATTGAACTT  | TACAATGTTA | TGTGTGATCT | 1600 |
|    | CCTGGGATTG  | AAGCCAGCTC  | CTAATAATGG | GACCCATGGA | 1640 |
|    | AGTTTGAATC  | ATCTCCTGCG  | CACTAATACC | TTCAGGCCAA | 1680 |
|    | CCATTGCCAGA | GGAAGTTACC  | AGACCCAATT | ATCCAGGGAT | 1720 |
|    | TATGTACCTT  | CAGTCTGATT  | TTGACCTGGG | CTGCACTTGT | 1760 |
|    | GATGATAAGG  | TAGAGCCAAA  | GAACAAGTTG | GATGAACTCA | 1800 |
|    | ACAAACGGCT  | TCATACAAAA  | GGGTCTACAG | AAGAGAGACA | 1840 |
|    | CCTCCTCTAT  | GGGCGACCTG  | CAGTGCTTTA | TCGGACTAGA | 1880 |
| 20 | TATGATGTCT  | TATATCACAC  | TGACTTTGAA | AGTGGTTATA | 1920 |
|    | GTGAAATATT  | CCTAATGCCA  | CTCTGGACAT | CATATACTGT | 1960 |
|    | TTCCAAACAG  | GCTGAGGTTT  | CCAGCGTTCC | TGACCATCTG | 2000 |
|    | ACCAGTTGCG  | TCCGGCCTGA  | TGTCCGTGTT | TCTCCGAGTT | 2040 |
|    | TCAGTCAGAA  | CTGTTTGGCC  | TACAAAAATG | ATAAGCAGAT | 2080 |
|    | GTCCACGGTA  | TTCTCTTTTC  | CTCCTTATCT | GAGCTCTTCA | 2120 |
|    | CCAGAGGCTA  | AATATGATGC  | ATTCTTTGTA | ACCAATATGG | 2160 |
| 25 | TTCCAATGTA  | TCCTGCTTTC  | AAACGGGTCT | GGAATTATTT | 2200 |
|    | CCAAAGGGTA  | TTGGTGAAGA  | AATATGCTTC | GGAAAGAAAT | 2240 |
|    | GGAGTTAACG  | TGATAAGTGG  | ACCAATCTTC | GACTATGACT | 2280 |
|    | ATGATGGCTT  | ACATGACACA  | GAAGACAAAA | TAAAACAGTA | 2320 |
|    | CGTGGAAGGC  | AGTTCCATTC  | CTGTTCCAAC | TCACTACTAC | 2360 |
|    | AGCATCATCA  | CCAGCTGTCT  | GGATTTCACT | CAGCCTGCCG | 2400 |
|    | ACAAGTGTGA  | CGGCCCTCTC  | TCTGTGTCTT | CCTTCATCCT | 2440 |
|    | CCGTCACCGG  | CCTGACAACG  | AGGAGAGCTG | CAATAGCTCA | 2480 |
| 30 | GAGGACGAAT  | CAAAATGGGT  | AGAAGAACTC | ATGAAGATGC | 2520 |
|    | ACACGGCTAG  | GGTGCGTGAC  | ATTGAACATC | TCACCAGCCT | 2560 |
|    | GGACTTCTTC  | CGAAAGACCA  | GCCGCAGCTA | CCCAGAAATC | 2600 |
|    | CTGACACTCA  | AGACATACCT  | GCATACATAT | GAGAGCGAGA | 2640 |
|    | TTTAACTTTT  | TGAGCATCTG  | CAGTACAGTC | TTATCAACTG | 2680 |
|    | GTTGTATATT  | TTTATATTGT  | TTTTGTATTT | ATTAATTTGA | 2720 |
|    | AACCAGGACA  | TTAAAAATGT  | TAGTATTTTA | ATCCTGTACC | 2760 |
|    | AAATCTGACA  | TATTATGCCT  | GAATGACTCC | ACTGTTTTTC | 2800 |
| 35 | TCTAATGCTT  | GATTTAGGTA  | GCCTTGTGTT | CTGAGTAGAG | 2840 |

- 86 -

CTTGTAATAA ATACTGCAGC TTGAGTTTTT AGTGGAAGCT 2880  
 TCTAAATGGT GCTGCAGATT TGATATTTGC ATTGAGGAAA 2920  
 TATTAATTTT CCAATGCACA GTTGCCACAT TTAGTCCTGT 2960  
 ACTGTATGGA AACACTGATT TTGTAAAGTT GCCTTTATTT 3000  
 GCTGTTAACT GTTAACTATG ACAGATATAT TTAAGCCTTA 3040  
 TAAACCAATC TTAAACATAA TAAATCACAC ATTCAGTTTT 3080  
 5 TTCTGGTAAA AAAAAAAAAA AAAA 3104

## (2) INFORMATION FOR SEQ ID NO:67:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 861
  - (B) TYPE: amino acid
  - 10 (C) STRANDEDNESS: single
  - (D) TOPOLOGY: Unknown
- (ii) MOLECULE TYPE: protein
- (iii) HYPOTHETICAL: No
- 15 (ix) FEATURE:
- (A) NAME/KEY: N-tera 2D1 ATX protein
  - (B) LOCATION:
  - (C) IDENTIFICATION METHOD:
  - (D) OTHER INFORMATION:

## 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

Met Ala Arg Arg Ser Ser Phe Gln Ser Cys Gln Ile  
 1 5 10  
 Ile Ser Leu Phe Thr Phe Ala Val Gly Val Asn Ile  
 15 20  
 Cys Leu Gly Phe Thr Ala His Arg Ile Lys Arg Ala  
 25 30 35  
 25 Glu Gly Trp Glu Glu Gly Pro Pro Thr Val Leu Ser  
 40 45  
 Asp Ser Pro Trp Thr Asn Ile Ser Gly Ser Cys Lys  
 50 55 60  
 Gly Arg Cys Phe Glu Leu Gln Glu Ala Gly Pro Pro  
 65 70  
 Asp Cys Arg Cys Asp Asn Leu Cys Lys Ser Tyr Thr  
 75 80  
 30 Ser Cys Cys His Asp Phe Asp Glu Leu Cys Leu Lys  
 85 90 95  
 Thr Ala Arg Ala Trp Glu Cys Thr Lys Asp Arg Cys  
 100 105  
 Gly Glu Val Arg Asn Glu Glu Asn Ala Cys His Cys  
 110 115 120  
 Ser Glu Asp Cys Leu Ala Arg Gly Asp Cys Cys Thr  
 125 130  
 35

- 87 -

|    |     |     |     |     |     |     |     |     |     |     |     |     |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| °  | Asn | Tyr | Gln | Val | Val | Cys | Lys | Gly | Glu | Ser | His | Trp |
|    |     |     | 135 |     |     |     |     | 140 |     |     |     |     |
|    | Val | Asp | Asp | Asp | Cys | Glu | Glu | Ile | Lys | Ala | Ala | Glu |
|    | 145 |     |     |     |     | 150 |     |     |     | 155 |     |     |
|    | Cys | Pro | Ala | Gly | Phe | Val | Arg | Pro | Pro | Leu | Ile | Ile |
|    |     |     | 160 |     |     |     |     |     | 165 |     |     |     |
|    | Phe | Ser | Val | Asp | Gly | Phe | Arg | Ala | Ser | Tyr | Met | Lys |
|    | 170 |     |     |     |     |     | 175 |     |     |     |     | 180 |
| 5  | Lys | Gly | Ser | Lys | Val | Met | Pro | Asn | Ile | Glu | Lys | Leu |
|    |     |     |     | 185 |     |     |     |     |     | 190 |     |     |
|    | Arg | Ser | Cys | Gly | Thr | His | Ser | Pro | His | Met | Arg | Pro |
|    |     |     | 195 |     |     |     |     | 200 |     |     |     |     |
|    | Val | Tyr | Pro | Thr | Lys | Thr | Phe | Pro | Asn | Leu | Tyr | Thr |
|    | 205 |     |     |     |     | 210 |     |     |     |     | 215 |     |
| 10 | Leu | Ala | Thr | Gly | Leu | Tyr | Pro | Glu | Ser | His | Gly | Ile |
|    |     |     | 220 |     |     |     |     |     | 225 |     |     |     |
|    | Val | Gly | Asn | Ser | Met | Tyr | Asp | Pro | Val | Phe | Asp | Ala |
|    | 230 |     |     |     |     |     | 235 |     |     |     |     | 240 |
|    | Thr | Phe | His | Leu | Arg | Gly | Arg | Glu | Lys | Phe | Asn | His |
|    |     |     |     | 245 |     |     |     |     |     | 250 |     |     |
|    | Arg | Trp | Trp | Gly | Gly | Gln | Pro | Leu | Trp | Ile | Thr | Ala |
|    |     |     | 255 |     |     |     |     | 260 |     |     |     |     |
| 15 | Thr | Lys | Gln | Arg | Gly | Glu | Ser | Trp | Asn | Ile | Leu | Leu |
|    | 265 |     |     |     |     | 270 |     |     |     |     | 275 |     |
|    | Val | Cys | Cys | His | Pro | Ser | Arg | Ala | Glu | Ile | Leu | Thr |
|    |     |     |     | 280 |     |     |     |     | 285 |     |     |     |
|    | Ile | Leu | Gln | Trp | Leu | Thr | Leu | Pro | Asp | His | Glu | Arg |
|    | 290 |     |     |     |     |     | 295 |     |     |     |     | 300 |
|    | Leu | Arg | Ser | Met | Pro | Ser | Ile | Leu | Ser | Asn | Leu | Ile |
|    |     |     |     | 305 |     |     |     |     |     | 310 |     |     |
| 20 | Ser | Leu | Asp | Thr | Asn | Met | Pro | Phe | Gly | Pro | Glu | Met |
|    |     |     | 315 |     |     |     |     | 320 |     |     |     |     |
|    | Thr | Asn | Pro | Leu | Arg | Glu | Ile | Asp | Lys | Ile | Val | Gly |
|    | 325 |     |     |     |     | 330 |     |     |     |     | 335 |     |
|    | Gln | Leu | Met | Asp | Gly | Leu | Lys | Gln | Leu | Lys | Leu | His |
|    |     |     | 340 |     |     |     |     |     | 345 |     |     |     |
|    | Arg | Cys | Val | Asn | Val | Ile | Phe | Val | Gly | Asp | His | Gly |
|    | 350 |     |     |     |     |     | 355 |     |     |     |     | 360 |
| 25 | Met | Glu | Asp | Val | Thr | Cys | Asp | Arg | Thr | Glu | Phe | Leu |
|    |     |     |     | 365 |     |     |     |     |     | 370 |     |     |
|    | Ser | Asn | Tyr | Leu | Thr | Asn | Val | Asp | Asp | Ile | Thr | Leu |
|    |     |     | 375 |     |     |     |     | 380 |     |     |     |     |
|    | Val | Pro | Gly | Thr | Leu | Gly | Ile | Arg | Ser | Lys | Phe | Ser |
|    | 385 |     |     |     |     | 390 |     |     |     |     | 395 |     |
|    | Asn | Asn | Ala | Lys | Tyr | Asp | Pro | Lys | Ala | Ile | Ile | Ala |
|    |     |     | 400 |     |     |     |     |     | 405 |     |     |     |
| 30 | Asn | Leu | Thr | Cys | Lys | Lys | Pro | Asp | Gln | His | Phe | Lys |
|    | 410 |     |     |     |     |     | 415 |     |     |     |     | 420 |
|    | Pro | Tyr | Leu | Lys | Gln | His | Leu | Pro | Lys | Arg | Leu | His |
|    |     |     |     | 425 |     |     |     |     |     | 430 |     |     |
|    | Tyr | Ala | Asn | Asn | Arg | Arg | Ile | Glu | Asp | Ile | His | Leu |
|    |     |     | 435 |     |     |     |     | 440 |     |     |     |     |
| 35 | Leu | Val | Glu | Arg | Arg | Trp | His | Val | Ala | Arg | Lys | Pro |
|    | 445 |     |     |     |     | 450 |     |     |     |     | 455 |     |

- 88 -

Leu Asp Val Tyr Lys Lys Pro Ser Gly Lys Cys Phe  
                             460                            465  
 Phe Gln Gly Asp His Gly Phe Asp Asn Lys Val Asn  
           470  475                            480  
 Ser Met Gln Thr Val Phe Val Gly Tyr Gly Pro Thr  
                             485                            490  
 5 Phe Lys Tyr Lys Thr Lys Val Pro Pro Phe Glu Asn  
                             495                            500  
 Ile Glu Leu Tyr Asn Val Met Cys Asp Leu Leu Gly  
 505  510                            515  
 Leu Lys Pro Ala Pro Asn Asn Gly Thr His Gly Ser  
                             520                            525  
 Leu Asn His Leu Leu Arg Thr Asn Thr Phe Arg Pro  
           530                            535                            540  
 10 Thr Met Pro Glu Glu Val Thr Arg Pro Asn Tyr Pro  
                             545                            550  
 Gly Ile Met Tyr Leu Gln Ser Asp Phe Asp Leu Gly  
                             555                            560  
 Cys Thr Cys Asp Asp Lys Val Glu Pro Lys Asn Lys  
 656  570                            575  
 Leu Asp Glu Leu Asn Lys Arg Leu His Thr Lys Gly  
                             580                            585  
 15 Ser Thr Glu Glu Arg His Leu Leu Tyr Gly Arg Pro  
           590                            595                            600  
 Ala Val Leu Tyr Arg Thr Arg Tyr Asp Val Leu Tyr  
                             605                            610  
 His Thr Asp Phe Glu Ser Gly Tyr Ser Glu Ile Phe  
                             615                            620  
 Leu Met Pro Leu Trp Thr Ser Tyr Thr Val Ser Lys  
 625  630                            635  
 20 Gln Ala Glu Val Ser Ser Val Pro Asp His Leu Thr  
                             640                            645  
 Ser Cys Val Arg Pro Asp Val Arg Val Ser Pro Ser  
           650                            655                            660  
 Phe Ser Gln Asn Cys Leu Ala Tyr Lys Asn Asp Lys  
                             665                            670  
 25 Gln Met Ser Tyr Gly Phe Leu Phe Pro Pro Tyr Leu  
                             675                            680  
 Ser Ser Ser Pro Glu Ala Lys Tyr Asp Ala Phe Leu  
 685  690                            695  
 Val Thr Asn Met Val Pro Met Tyr Pro Ala Phe Lys  
                             700                            705  
 Arg Val Trp Asn Tyr Phe Gln Arg Val Leu Val Lys  
           710                            715                            720  
 30 Lys Tyr Ala Ser Glu Arg Asn Gly Val Asn Val Ile  
                             725                            730  
 Ser Gly Pro Ile Phe Asp Tyr Asp Tyr Asp Gly Leu  
                             735                            740  
 His Asp Thr Glu Asp Lys Ile Lys Gln Tyr Val Glu  
 745  750                            755  
 Gly Ser Ser Ile Pro Val Pro Thr His Tyr Tyr Ser  
                             760                            765  
 35

- 89 -

° Ile Ile Thr Ser Cys Leu Asp Phe Thr Gln Pro Ala  
       770                              775                              780  
 Asp Lys Cys Asp Gly Pro Leu Ser Val Ser Ser Phe  
                               785                              790  
 Ile Leu Arg His Arg Pro Asp Asn Glu Glu Ser Cys  
                               795                              800  
 Asn Ser Ser Glu Asp Glu Ser Lys Trp Val Glu Glu  
 5 805                              810                              815  
 Leu Met Lys Met His Thr Ala Arg Val Arg Asp Ile  
                               820                              825  
 Glu His Leu Thr Ser Leu Asp Phe Phe Arg Lys Thr  
       830                              835                              840  
 Ser Arg Ser Tyr Pro Glu Ile Leu Thr Leu Lys Thr  
                               845                              850  
 10 Tyr Leu His Thr Tyr Glu Ser Glu Ile  
                               855                              860

## (2) INFORMATION FOR SEQ ID NO:68:

(i) SEQUENCE CHARACTERISTICS:  
       (A) LENGTH: 3251  
       (B) TYPE: nucleic acid  
 15 (C) STRANDEDNESS: double  
       (D) TOPOLOGY: Unknown  
  
 (ii) MOLECULE TYPE: cDNA  
  
 (iii) HYPOTHETICAL: No  
  
 (ix) FEATURE:  
 20 (A) NAME/KEY: A2058 ATX cDNA  
       (B) LOCATION:  
       (C) IDENTIFICATION METHOD:  
       (D) OTHER INFORMATION:  
  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

|    |              |             |            |            |     |
|----|--------------|-------------|------------|------------|-----|
| 25 | CGTGAAGGCA   | AAGAGAACAC  | GCTGCAAAAG | GCTTCCAAGA | 40  |
|    | ATCCTCGACA   | TGGCAAGGAG  | GAGCTCGTTC | CAGTCGTGTC | 80  |
|    | AGATAATATC   | CCTGTTCACT  | TTTGCCGTTG | GAGTCAGTAT | 120 |
|    | CTGCTTAGGA   | TTCAC TGCAC | ATCGAATTAA | GAGAGCAGAA | 160 |
|    | GGATGGGAGG   | AAGGTCCTCC  | TACAGTGCTA | TCAGACTCCC | 200 |
|    | CCTGGACCAA   | CATCTCCGGA  | TCTTGCAAGG | GCAGGTGCTT | 240 |
|    | TGAACTTCAA   | GAGGCTGGAC  | CTCCTGATTG | TCGCTGTGAC | 280 |
| 30 | AACTTGTGTA   | AGAGCTATAC  | CAGTTGCTGC | CATGACTTTG | 320 |
|    | ATGAGCTGTG   | TTTGAAGACA  | GCCCGTGGCT | GGGAGTGTAC | 360 |
|    | TAAGGACAGA   | TGTGGAGAAG  | TCAGAAATGA | AGAAAATGCC | 400 |
|    | TGTCAC T GCT | CAGAGGACTG  | CTTGGCCAGG | GGAGACTGCT | 440 |
|    | GTACCAATTA   | CCAAGTGGTT  | TGCAAAGGAG | AGTCGCATTG | 480 |
|    | GGTTGATGAT   | GACTGTGAGG  | AAATAAAGGC | CGCAGAATGC | 520 |
|    | CCTGCAGGGT   | TTGTTGCCCC  | TCCATTAATC | ATCTTCTCCG | 560 |
|    | TGGATGGCTT   | CCGTGCATCA  | TACATGAAGA | AAGGCAGCAA | 600 |
| 35 | AGTCATGCCT   | AATATTGAAA  | AACTAAGGTC | TTGTGGCACA | 640 |

- 90 -

|    |             |             |            |             |      |
|----|-------------|-------------|------------|-------------|------|
|    | CACTCTCCCT  | ACATGAGGCC  | GGTGTACCCA | ACTAAAACCT  | 680  |
|    | TTCCTAACTT  | ATACACTTTG  | GCCACTGGGC | TATATCCAGA  | 720  |
|    | ATCACATGGA  | ATTGTTGGCA  | ATTCAATGTA | TGATCCTGTA  | 760  |
|    | TTTGATGCCA  | CTTTTCATCT  | GCGAGGGCGA | GAGAAATTTA  | 800  |
|    | ATCATAGATG  | GTGGGGAGGT  | CAACCGCTAT | GGATTACAGC  | 840  |
|    | CACCAAGCAA  | GGGGTGAAAG  | CTGGAACATT | CTTTTGGTCT  | 880  |
| 5  | GTTGTTCATCC | CTCACGAGCG  | GAGAATATTA | ACCATATTGC  | 920  |
|    | GGTGGCTCAC  | CCTGCCAGAT  | CATGAGAGGC | CTTCGGTCTA  | 960  |
|    | TGCCCTTCTAT | TCTGAGCAAC  | CTGATTTCTC | TGGACACAAA  | 1000 |
|    | TATGGCCCTT  | TCGGCCCTGA  | GGAGAGTAGT | TATGGCTCAC  | 1040 |
|    | CTTTTACTCC  | GGCTAAGAGA  | CCTAAGAGGA | AAGTTGCCCC  | 1080 |
|    | TAAGAGGAGA  | CAGGAAAGAC  | CAGTTGCTCC | TCCAAAGAAA  | 1120 |
|    | AGAAGAAGAA  | AAATACATAG  | GATGGATCAT | TATGCTGCGG  | 1160 |
|    | AAACTCGTCA  | GGACAAAATG  | ACAAATCCTC | TGAGGGAAAT  | 1200 |
| 10 | CGACAAAATT  | GTGGGGCAAT  | TAATGGATGG | ACTGAAACAA  | 1240 |
|    | CTAAAACCTGC | GTCGGTGTGT  | CAACGTCATC | TTTGTCTGGAG | 1280 |
|    | ACCATGGAAT  | GGAAGATGTC  | ACATGTGATA | GAAGTGAAGT  | 1320 |
|    | CTTGAGTAAT  | TACCTAACTA  | ATGTGGATGA | TATTACTTTA  | 1360 |
|    | GTGCCTGGAA  | CTCTAGGAAG  | AATTCGATCC | AAATTTAGCA  | 1400 |
|    | ACAATGCTAA  | ATATGACCCC  | AAAGCCATTA | TTGCCAATCT  | 1440 |
|    | CACGTGTAAA  | AAACCAGATC  | AGCACTTTAA | GCCTTACTTG  | 1480 |
| 15 | AAACAGCACC  | TTCCCAAACG  | TTTGCACTAT | GCCAACAACA  | 1520 |
|    | GAAGAATTGA  | GGATATCCAT  | TTATTGGTGG | AACGCAGATG  | 1560 |
|    | GCATGTTGCA  | AGGAAACCTT  | TGGATGTTTA | TAAGAAACCA  | 1600 |
|    | TCAGGAAAAT  | GCTTTTTCCTA | GGGAGACCAC | GGATTTGATA  | 1640 |
|    | ACAAGGTCAA  | CAGCATGCAG  | ACTGTTTTTG | TAGGTTATGG  | 1680 |
|    | CCCAACATTT  | AAGTACAAGA  | CTAAAGTGCC | TCCATTTGAA  | 1720 |
|    | AACATTGAAC  | TTTACAATGT  | TATGTGTGAT | CTCCTGGGAT  | 1760 |
|    | TGAAGCCAGC  | TCCTAATAAT  | GGGACCCATG | GAAGTTTGAA  | 1800 |
| 20 | TCATCTCCTG  | CGCACTAATA  | CCTTCAGGCC | AACCATGCCA  | 1840 |
|    | GAGGAACTTA  | CCAGACCCAA  | TTATCCAGGG | ATTATGTACC  | 1880 |
|    | TTCAGTCTGA  | TTTTGACCTG  | GGCTGCACCT | GTGATGATAA  | 1920 |
|    | GGTAGAGCCA  | AAGAACAAGT  | TGGATGAACT | CAACAAACGG  | 1960 |
|    | CTTCATACAA  | AAGGGTCTAC  | AGAAGAGAGA | CACCTCCTCT  | 2000 |
|    | ATGGGCGACC  | TGCAGTGCTT  | TATCGGACTA | GATATGATAT  | 2040 |
|    | CTTATATCAC  | ACTGACTTTG  | AAAGTGGTTA | TAGTGAAATA  | 2080 |
| 25 | TTCTTAATGC  | TACTCTGGAC  | ATCATATACT | GTTTCCAAAC  | 2120 |
|    | AGGCTGAGGT  | TTCCAGCGTT  | CCTGACCATC | TGACCAGTTG  | 2160 |
|    | CGTCCGGCCT  | GATGTCCGTG  | TTTCTCCGAG | TTTCAGTCAG  | 2200 |
|    | AACTGTTTGG  | CCTACAAAAA  | TGATAAGCAG | ATGTCCTACG  | 2240 |
|    | GATTCCCTCT  | TCCTCCTTAT  | CTGAGCTCTT | CACCAGAGGC  | 2280 |
|    | TAAATATGAT  | GCATTCCCTG  | TAACCAATAT | GGTTCCAATG  | 2320 |
|    | TATCCTGCTT  | TCAAACGGGT  | CTGGAATTAT | TTCCAAAGGG  | 2360 |
| 30 | TATTGGTGAA  | GAAATATGCT  | TCGGAAAGAA | ATGGAGTTAA  | 2400 |
|    | CGTGATAAGT  | GGACCAATCT  | TCGACTATGA | CTATGATGGC  | 2440 |
|    | TTACATGACA  | CAGAAGACAA  | AATAAAACAG | TACGTGGAAG  | 2480 |
|    | GCAGTTCCAT  | TCCTGTTCCA  | ACTCACTACT | ACAGCATCAT  | 2520 |
|    | CACCAGCTGT  | CTGGATTTC   | CTCAGCCTGC | CGACAAGTGT  | 2560 |
|    | GACGGCCCTC  | TCTCTGTGTC  | CTCCTTCATC | CTGCCTCACC  | 2600 |
|    | GGCCTGACAA  | CGAGGAGAGC  | TGCAATAGCT | CAGAGGACGA  | 2640 |
|    | ATCAAAATGG  | GTAAGAAGAC  | TCATGAAGAT | GCACACAGCT  | 2680 |
| 35 | AGGGTGCGTG  | ACATTGAACA  | TCTCACCAGC | CTGGACTTCT  | 2720 |
|    | TCCGAAAGAC  | CAGCCGCAGC  | TACCCAGAAA | TCCTGACACT  | 2760 |



- 91 -

CAAGACATAC CTGCATACAT ATGAGAGCGA GATTTAACTT 2800  
 TCTGAGCATC TGCAGTACAG TCTTATCAAC TGGTTGTATA 2840  
 TTTTATATATT GTTTTGTAT TTATTAATTT GAAACCAGGA 2880  
 CATTAAAAAT GTTAGTATTT TAATCCTGTA CCAAATCTGA 2920  
 CATATTATGC CTGAATGACT CCACTGTTTT TCTCTAATGC 2960  
 TTGATTTAGG TAGCCTTGTG TTCTGAGTAG AGCTTGTAAT 3000  
 5 AAATACTGCA GCTTGAGAAA AAGTGGAAGC TTCTAAATGG 3040  
 TGCTGCAGAT TTGATATTTG CATTGAGGAA ATATTAATTT 3080  
 TCCAATGCAC AGTTGCCACA TTTAGTCCTG TACTGTATGG 3120  
 AAACACTGAT TTTGTAAAGT TGCCTTTATT TGCTGTTAAC 3160  
 TGTTAACTAT GACAGATATA TTAAAGCCTT ATAAACCAAT 3200  
 CTTAAACATA ATAAATCACA CATTCAGTTT TAAAAAATAA 3240  
 AAAAAAAAAA A 3251

10

(2) INFORMATION FOR SEQ ID NO:69:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 915  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: Unknown

15

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: No

(ix) FEATURE:

- (A) NAME/KEY: A2058 ATX protein  
 (B) LOCATION:  
 (C) IDENTIFICATION METHOD:  
 (D) OTHER INFORMATION:

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

25 Met Ala Arg Arg Ser Ser Phe Gln Ser Cys Gln Ile  
     1                    5                    10  
 Ile Ser Leu Phe Thr Phe Ala Val Gly Val Ser Ile  
           15                    20  
 Cys Leu Gly Phe Thr Ala His Arg Ile Lys Arg Ala  
   25                    30                    35  
 Glu Gly Trp Glu Glu Gly Pro Pro Thr Val Leu Ser  
           40                    45  
 30 Asp Ser Pro Trp Thr Asn Ile Ser Gly Ser Cys Lys  
     50                    55                    60  
 Gly Arg Cys Phe Glu Leu Gln Glu Ala Gly Pro Pro  
           65                    70  
 Asp Cys Arg Cys Asp Asn Leu Cys Lys Ser Tyr Thr  
     75                    80  
 Ser Cys Cys His Asp Phe Asp Glu Leu Cys Leu Lys  
   85                    90                    95

35

- 92 -

° Thr Ala Arg Gly Trp Glu Cys Thr Lys Asp Arg Cys  
                                   100                                  105  
 Gly Glu Val Arg Asn Glu Glu Asn Ala Cys His Cys  
           110                                  115                                  120  
 Ser Glu Asp Cys Leu Ala Arg Gly Asp Cys Cys Thr  
                                   125                                  130  
 Asn Tyr Gln Val Val Cys Lys Gly Glu Ser His Trp  
                                   135                                  140  
 5 Val Asp Asp Asp Cys Glu Glu Ile Lys Ala Ala Glu  
    145                                  150                                  155  
 Cys Pro Ala Gly Phe Val Arg Pro Pro Leu Ile Ile  
                                   160                                  165  
 Phe Ser Val Asp Gly Phe Arg Ala Ser Tyr Met Lys  
           170                                  175                                  180  
 10 Lys Gly Ser Lys Val Met Pro Asn Ile Glu Lys Leu  
                                   185                                  190  
 Arg Ser Cys Gly Thr His Ser Pro Tyr Met Arg Pro  
                                   195                                  200  
 Val Tyr Pro Thr Lys Thr Phe Pro Asn Leu Tyr Thr  
    205                                  210                                  215  
 Leu Ala Thr Gly Leu Tyr Pro Glu Ser His Gly Ile  
                                   220                                  225  
 15 Val Gly Asn Ser Met Tyr Asp Pro Val Phe Asp Ala  
    230                                  235                                  240  
 Thr Phe His Leu Arg Gly Arg Glu Lys Phe Asn His  
                                   245                                  250  
 Arg Trp Trp Gly Gly Gln Pro Leu Trp Ile Thr Ala  
                                   255                                  260  
 Thr Lys Gln Gly Val Lys Ala Gly Thr Phe Phe Trp  
    265                                  270                                  275  
 20 Ser Val Val Ile Pro His Glu Arg Arg Ile Leu Thr  
                                   280                                  285  
 Ile Leu Arg Trp Leu Thr Leu Pro Asp His Glu Arg  
    290                                  295                                  300  
 Pro Ser Val Tyr Ala Phe Tyr Ser Glu Gln Pro Asp  
                                   305                                  310  
 Phe Ser Gly His Lys Tyr Gly Pro Phe Gly Pro Glu  
                                   315                                  320  
 25 Glu Ser Ser Tyr Gly Ser Pro Phe Thr Pro Ala Lys  
    325                                  330                                  335  
 Arg Pro Lys Arg Lys Val Ala Pro Lys Arg Arg Gln  
                                   340                                  345  
 Glu Arg Pro Val Ala Pro Pro Lys Lys Arg Arg Arg  
    350                                  355                                  360  
 Lys Ile His Arg Met Asp His Tyr Ala Ala Glu Thr  
                                   365                                  370  
 30 Arg Gln Asp Lys Met Thr Asn Pro Leu Arg Glu Ile  
                                   375                                  380  
 Asp Lys Ile Val Gly Gln Leu Met Asp Gly Leu Lys  
    385                                  390                                  395  
 Gln Leu Lys Leu Arg Arg Cys Val Asn Val Ile Phe  
                                   400                                  405  
 35 Val Gly Asp His Gly Met Glu Asp Val Thr Cys Asp  
    410                                  415                                  420

- 93 -

Arg Thr Glu Phe Leu Ser Asn Tyr Leu Thr Asn Val  
 425 430  
 Asp Asp Ile Thr Leu Val Pro Gly Thr Leu Gly Arg  
 435 440  
 Ile Arg Ser Lys Phe Ser Asn Asn Ala Lys Tyr Asp  
 445 450 455  
 5 Pro Lys Ala Ile Ile Ala Asn Leu Thr Cys Lys Lys  
 460 465  
 Pro Asp Gln His Phe Lys Pro Tyr Leu Lys Gln His  
 470 475 480  
 Leu Pro Lys Arg Leu His Tyr Ala Asn Asn Arg Arg  
 485 490  
 Ile Glu Asp Ile His Leu Leu Val Glu Arg Arg Trp  
 495 500  
 10 His Val Ala Arg Lys Pro Leu Asp Val Tyr Lys Lys  
 505 510 515  
 Pro Ser Gly Lys Cys Phe Phe Gln Gly Asp His Gly  
 520 525  
 Phe Asp Asn Lys Val Asn Ser Met Gln Thr Val Phe  
 530 535 540  
 Val Gly Tyr Gly Pro Thr Phe Lys Tyr Lys Thr Lys  
 545 550  
 15 Val Pro Pro Phe Glu Asn Ile Glu Leu Tyr Asn Val  
 555 560  
 Met Cys Asp Leu Leu Gly Leu Lys Pro Ala Pro Asn  
 565 570 575  
 Asn Gly Thr His Gly Ser Leu Asn His Leu Leu Arg  
 580 585  
 Thr Asn Thr Phe Arg Pro Thr Met Pro Glu Glu Val  
 590 595 600  
 20 Thr Arg Pro Asn Tyr Pro Gly Ile Met Tyr Leu Gln  
 605 610  
 Ser Asp Phe Asp Leu Gly Cys Thr Cys Asp Asp Lys  
 615 620  
 Val Glu Pro Lys Asn Lys Leu Asp Glu Leu Asn Lys  
 625 630 635  
 25 Arg Leu His Thr Lys Gly Ser Thr Glu Glu Arg His  
 640 645  
 Leu Leu Tyr Gly Arg Pro Ala Val Leu Tyr Arg Thr  
 650 655 660  
 Arg Tyr Asp Ile Leu Tyr His Thr Asp Phe Glu Ser  
 665 670  
 Gly Tyr Ser Glu Ile Phe Leu Met Leu Leu Trp Thr  
 675 680  
 30 Ser Tyr Thr Val Ser Lys Gln Ala Glu Val Ser Ser  
 685 690 695  
 Val Pro Asp His Leu Thr Ser Cys Val Arg Pro Asp  
 700 705  
 Val Arg Val Ser Pro Ser Phe Ser Gln Asn Cys Leu  
 710 715 720  
 Ala Tyr Lys Asn Asp Lys Gln Met Ser Tyr Gly Phe  
 725 730  
 35

- 94 -

|    |  |     |     |            |            |            |            |            |            |            |            |            |            |
|----|--|-----|-----|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| °  |  | Leu | Phe | Pro<br>735 | Pro        | Tyr        | Leu        | Ser        | Ser<br>740 | Ser        | Pro        | Glu        | Ala        |
|    |  | Lys | Tyr | Asp        | Ala        | Phe        | Leu<br>750 | Val        | Thr        | Asn        | Met        | Val<br>755 | Pro        |
|    |  | Met | Tyr | Pro        | Ala<br>760 | Phe        | Lys        | Arg        | Val        | Trp<br>765 | Asn        | Tyr        | Phe        |
|    |  | Gln | Arg | Val        | Leu        | Val        | Lys<br>775 | Lys        | Tyr        | Ala        | Ser        | Glu        | Arg<br>780 |
| 5  |  | Asn | Gly | Val        | Asn        | Val<br>785 | Ile        | Ser        | Gly        | Pro        | Ile<br>790 | Phe        | Asp        |
|    |  | Tyr | Asp | Tyr<br>795 | Asp        | Gly        | Leu        | His        | Asp<br>800 | Thr        | Glu        | Asp        | Lys        |
|    |  | Ile | Lys | Gln        | Tyr        | Val        | Glu<br>810 | Gly        | Ser        | Ser        | Ile        | Pro<br>815 | Val        |
| 10 |  | Pro | Thr | His<br>820 | Tyr        | Tyr        | Ser        | Ile        | Ile        | Thr<br>825 | Ser        | Cys        | Leu        |
|    |  | Asp | Phe | Thr        | Gln        | Pro        | Ala<br>835 | Asp        | Lys        | Cys        | Asp        | Gly        | Pro<br>840 |
|    |  | Leu | Ser | Val        | Ser        | Ser<br>845 | Phe        | Ile        | Leu        | Pro        | His<br>850 | Arg        | Pro        |
|    |  | Asp | Asn | Glu<br>855 | Glu        | Ser        | Cys        | Asn        | Ser<br>860 | Ser        | Glu        | Asp        | Glu        |
| 15 |  | Ser | Lys | Trp        | Val        | Glu        | Glu<br>870 | Leu        | Met        | Lys        | Met        | His<br>875 | Thr        |
|    |  | Ala | Arg | Val        | Arg<br>880 | Asp        | Ile        | Glu        | His        | Leu<br>885 | Thr        | Ser        | Leu        |
|    |  | Asp | Phe | Phe        | Arg        | Lys        | Thr        | Ser<br>895 | Arg        | Ser        | Tyr        | Pro        | Glu<br>900 |
|    |  | Ile | Leu | Thr        | Leu        | Lys<br>905 | Thr        | Tyr        | Leu        | His        | Thr<br>910 | Tyr        | Glu        |
| 20 |  | Ser | Glu | Ile<br>915 |            |            |            |            |            |            |            |            |            |

25

30

35

- 95 -

°  
CLAIMS:

1. A DNA segment coding for a polypeptide comprising an amino acid sequence corresponding to autotaxin, or a fragment thereof having at least 5 amino acids.

5           2. The DNA segment according to claim 1, wherein said DNA segment encodes the amino acid sequence selected from the group consisting of SEQ ID NO:1 through SEQ ID NO:11, SEQ ID NO:26 through SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38 through SEQ ID NO:52, SEQ ID NO:66  
10       and SEQ ID NO:69.

3. An isolated polypeptide comprising an amino acid sequence corresponding to autotaxin, or a fragment thereof having at least 5 amino acids.

15           4. The polypeptide according to claim 3, wherein said amino acid sequence comprises the amino acid sequence selected from the group consisting of SEQ ID NO:1 through SEQ ID NO:11, SEQ ID NO:26 through SEQ ID NO:34, SEQ ID NO:36, and SEQ ID NO:38 through SEQ ID NO:52, SEQ ID NO:67 and SEQ ID NO:69.

20           5. An isolated polypeptide bound to a solid support, comprising an amino acid sequence corresponding to autotaxin, or a fragment thereof having at least 5 amino acids.

25           6. The polypeptide according to claim 5, wherein said polypeptide comprises the amino acid sequence selected from the group consisting of the SEQ ID NO:1 through SEQ ID NO:11, SEQ ID NO:26 through SEQ ID NO:34, SEQ ID NO: 36, SEQ ID NO:38 through SEQ ID NO:52., SEQ ID NO:67 and SEQ ID NO:69.

30           7. A recombinant DNA molecule comprising a vector and the DNA segment according to claim 1.

8. A cell that contains the recombinant DNA molecule according to claim 7.

35           9. An antibody having binding affinity for autotaxin, or binding fragment thereof.

- 96 -

10. A method of producing a recombinant autotaxin polypeptide said method comprising:
- culturing a cell containing the recombinant DNA molecule of claim 7 under conditions such that the DNA segment is expressed, producing said polypeptide; and
  - isolating said polypeptide.
11. A method of purifying the autotaxin peptide of claim 3, comprising the steps of:
- i) collecting and concentrating supernatant from cultured A2058 human melanoma cells whereby a first preparation of said peptide is produced;
  - ii) salt fractionating said first preparation to produce a second peptide preparation;
  - iii) isolating said peptide from said second preparation so that said peptide is obtained in substantially pure form.
12. The method of claim 11, wherein said isolating step is effected by column chromatography.
13. An isolated DNA encoding an autotaxin protein or fragment thereof wherein said DNA includes a nucleic acid sequence selected from the group consisting of SEQ ID NO:35, SEQ ID NO:37 and SEQ ID NO:38.
14. The DNA segment according to claim 1, wherein said DNA fragment comprises any one of the SEQ ID NO:12 through SEQ ID NO:25, or SEQ ID NO:39 through SEQ ID NO:52.
15. The DNA segment according to claim 13 wherein said DNA segment comprises any one of the SEQ ID NO:12 through SEQ ID NO:25.
16. An isolated polypeptide comprising an amino acid sequence corresponding to autotaxin.
17. A polypeptide bound to a solid support and comprising an amino acid sequence corresponding to autotaxin.
18. A recombinant autotaxin polypeptide

- 97 -

° according to claim 3.

19. An isolated polypeptide according to claim 3 having cell motility activity.

5

10

15



20

25

30



35

1/15



**FIG. 1****FIG. 2**




2/15

**FIG. 3****FIG. 4**

3/15

**FIG. 5****FIG. 8**

4/15

**FIG. 6A**

5/15

FIG. 6C








FIG. 6B



6/15


**FIG. 7****FIG. 16**

7/15


**FIG. 9****FIG. 10**

|             |       | Upper Walls |            |            |
|-------------|-------|-------------|------------|------------|
|             |       | 0           | 0.01%      | 0.1%       |
| Lower Walls | 0     | 4.8 ± 0.3   | 13.7 ± 0.8 | 33.8 ± 1.6 |
|             | 0.01% | 45.4 ± 4.0  | 39.3 ± 2.6 | 34.9 ± 1.4 |
|             | 0.1%  | 75.6 ± 1.8  | 58.3 ± 3.1 | 41.0 ± 3.4 |

8/15

**FIG. 11**

9/15

**FIG. 12**



10/15

FIG. 13






FIG. 19




11/15

FIG. 14




12/15

FIG. 15



13/15

**FIG. 17A****FIG. 17B**

SUBSTITUTE SHEET (RULE 26)

14/15

## FIG. 18A

|      |  |     |
|------|--|-----|
| hATX | MARRSSFQSQIISLFTFAVGVSICLGTAAHRIKRAEGWEEGPTVLSDSPWNIISGCKGRCFELQACGPPDCRCDNLCKSYTSCCHDF              | 90  |
| hPC1 | MDVGEEPLEKAARARTAKDPNTYKVLISLVSVCVLTTIL.....GCIFG....LKPSCAKEVK.SCKGRCF...ERTFGNCRCDAAACVELGHCCLDY   | 84  |
| hATX | DELCLKTARGWECTKDRCGEVRNEENACHCEDCLARGDCCTNYQVCKGESHVDDDCCEIKAAECAGFVRPPLIIFSVDFGRASYMKKSKVMPNIE      | 190 |
| hPC1 | QETCIEPEHIWTCNFRGCEKRLTRSLCACDDCKDKGCCINYSVCQGEKSWVEEPCESINEPQCPAGFETPTLLFSLDGFRAEYLHTWGGLLPVIS      | 184 |
| hATX | KLRSCGTHSPYMRPVYPTKTFPNLYTLATGLYPESHGIVGSMYDPVFDFATFHLRGREKFNHRWGGQPLWITATKQGVKAGTFFWS               | 272 |
| hPC1 | KLKKGCTYTKMRPYPTKTFPNHYSIVTGLYPESHGIIDNKMYDPKMNASFSLKSKEKFNPEWKGEPIWVTAKYQGLKSGTFFWPGSDVEINGIFPDI    | 284 |
| hATX | .....VVIPHERIILTLRWLTPDHERPSVAFYSEQPDFSGHKYGPFGPESSYSPFTPAKHPKRKVA PKRQRPVAPPKRRRKIHRMDHYAAET        | 372 |
| hPC1 | YKMYNGSVPFEEIRILAVLQWLQPKDERPHFYTYLLEEDPSGHSYGPVSSE.....   | 336 |
| hATX | RQDKMTNPLREIDKIVQGLMDGLKQLKLRRCNVNIFVGHGMEDVTCRTEFLSNYLTNVDDITLVPGTLGRIR.SKFSNN.AKYDPKAIITANLTCKKPD  | 470 |
| hPC1 | .....VTKALQRVDGWGMLMDGLKELNLHRCNLILTSDHMEQGSCKKIYLNKYLGDVKNIKVIYGAARLPSDVPDKYYSNVEGIARNLSCREPN       | 432 |
| hATX | QHFKPYLKQHLPKRLHYANNRRIEDIHLVERRWHVARKPLDVYKPSGCKFFQDGHGFDNKNVSMQTVPVYGYGPTFKYKTKVPPFENIELYNVCMCDLIG | 570 |
| hPC1 | QHFKPYLKHLGPKRLHFAKSDRIEPLTFYLDPPQWLALNPSE..RKYCGSGF....HGSDNVFSNMQALFVGYGPGFKHGTADTFENIEVYNLMCDLLN  | 526 |

